

Measurements of the EMC Effect Using PVDIS Update

Seamus Riordan
Stony Brook University
`seamus.riordan@stonybrook.edu`

with

John Arrington
Argonne National Laboratory

Rakitha Beminiwattha
Syracuse University



September 11, 2015

- Motivation
- Proposed Experiment
- Anticipated Results and Systematics

- Went to PAC42 - deferred
 - Cited desire to run Hall C $^{48}\text{Ca}/^{40}\text{Ca}$ first
 - Concerns about beam time and radiation budget for site boundaries
- John Arrington has joined as spokesperson
- Working on updating proposal for resubmission
 - Need to consider strengthening (and clarifying) physics case
 - more detailed radiation calculations
 - ^{48}Ca target update with status

Modification Questions

- Possible new degrees of freedom must be considered for nucleon modification - can CSV effectively be induced in asymmetric nuclei?
- Are data such as those from NuTeV hinting at us that these effects are important?
- Is there a direct connection between nucleon and parton-level modification observables?



PVDIS proves new flavor combinations \rightarrow **isovector properties**

$$A_{PV} \sim \frac{\left| \begin{array}{c} \gamma^* \\ \text{---} \\ \text{---} \end{array} \right| \left| \begin{array}{c} Z^* \\ \text{---} \\ \text{---} \end{array} \right|}{\left| \begin{array}{c} \gamma^* \\ \text{---} \\ \text{---} \end{array} \right|^2} \sim 100 - 1000 \text{ ppm}$$

$$\approx -\frac{G_F Q^2}{4\sqrt{2}\pi\alpha} \left[a_1(x) + \frac{1 - (1-y)^2}{1 + (1-y)^2} a_3(x) \right], y = 1 - \frac{E'}{E}$$

$$a_1(x) = 2 \frac{\sum C_{1q} e_q (q + \bar{q})}{\sum e_q^2 (q + \bar{q})}, a_3(x) = 2 \frac{\sum C_{2q} e_q (q - \bar{q})}{\sum e_q^2 (q + \bar{q})}$$

Effective Weak Couplings

$$\begin{array}{ll} C_{1u} = -\frac{1}{2} + \frac{4}{3} \sin^2 \theta_W = -0.19 & C_{2u} = -\frac{1}{2} + 2 \sin^2 \theta_W = -0.03 \\ C_{1d} = \frac{1}{2} - \frac{2}{3} \sin^2 \theta_W = 0.34 & C_{2d} = \frac{1}{2} + 2 \sin^2 \theta_W = 0.03 \end{array}$$

PVDIS proves new flavor combinations \rightarrow **isovector properties**

$$A_{PV} \sim \frac{\left| \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \right|^2}{\left| \begin{array}{c} \text{Diagram 3} \end{array} \right|^2} \sim 100 - 1000 \text{ ppm}$$

$$\approx -\frac{G_F Q^2}{4\sqrt{2}\pi\alpha} \left[a_1(x) + \frac{1 - (1-y)^2}{1 + (1-y)^2} a_3(x) \right], y = 1 - \frac{E'}{E}$$

Symmetric nucleus limit

Probing *isovector* quantity through parity violation:


$$a_1 \simeq \frac{9}{5} - 4 \sin^2 \theta_W - \frac{12}{25} \left[\frac{u_A^+ - d_A^+}{u_A^+ + d_A^+} \right] + \dots$$

where $u_A = u$ in p and u in n

Nuclear Modification

- First observed in 1984 by EMC collaboration
- Showed reduced presence of partons in $0.3 < x < 0.7$
- Generally greater effect as one pushes to higher A
- Not due to simple binding effects - real modification of structure

General assumption of $u \leftrightarrow d$ for $p \leftrightarrow n$
PVDIS can test this




J. Gomez et al.,
PRD49 4348 (1994)

- Neutrino scattering (charged current and neutral current) is sensitive to different flavor combinations

Pachos-Wolfenstein relation:

$$R_{PW} \equiv \frac{\sigma(\nu_\mu N \rightarrow \nu_\mu X) - \sigma(\bar{\nu}_\mu N \rightarrow \bar{\nu}_\mu X)}{\sigma(\nu_\mu N \rightarrow \mu^- X) - \sigma(\bar{\nu}_\mu N \rightarrow \mu^+ X)} \sin^2 \theta_W(\mu)$$



$$= \lim_{\rightarrow \text{i.s.}} \frac{1}{2} - \sin^2 \theta_W$$



- Asymmetric nuclei (iron) need corrections
- CSV or IVEMC (effective CSV) will change this result and are **not well constrained by data**

Modeling - CBT Model



- Cloet et al. make predictions based on mean field calculations which give reasonable reproductions of SFs
- Explicit isovector terms are included constrained by symmetry energy
- Few percent effect in a_2 , larger at larger x



Cloet et al. PRL102 252301 (2009), Cloet et al. PRL109 182301 (2012)


Isvector Dependence? - SRC

- SRC show strong preference to n-p pairs over p-p pairs
- Show strong correlation to “plateau” parameter for $x > 1$ SFs
- Different high momentum tails for n and p in asymmetric nuclei could drive modification and IVEMC




Configuration

- Experimental configuration practically identical to approved SoLID PVDIS measurement
- Rates are better or comparable to existing LD₂ measurement



Target - ^{48}Ca

- ^{48}Ca target provides good balance between asymmetric target and not too high Z
- 12% radiator - photons and photoproduced pions are main background concerns



- Learned that ^{48}Ca is not isotopically pure and has $\sim 10\%$ ^{40}Ca and other contaminations - needs to be put under consideration
- Probably only going to have single target already in position - needs to be recast - holder needs to be redesigned

- Polarimetry and pions are main contributions
- Radiative working group has been established for PVDIS
- Dominant errors:

Effect	Uncertainty [%]
Polarimetry	0.4
Q^2 -dependent effects	0.2
Pions (bin-to-bin)	0.1-0.5
Radiative Corrections (bin-to-bin)	0.5-0.1
e^- from pair production	0.5-0.0
Total for any given bin	$\sim 0.5-0.8$

- Statistical uncertainty dominates

Radiation Summary

- Beam on target simulation using Remoll
 - Limited statistics – about 20% uncertainty on neutrons
- No Polythene yet

Radiation Power with No Poly

Type	E range (MeV)	Ca48	LD2	PREX_I	PREX_II
		Power (W/uA)	Power (W/uA)	Power (W/uA)	Power (W/uA)
e±	E<10	0.11	0.11	0.17	0.01
	10<E	0.18	0.16	1.29	0.06
Photons	E<10	0.02	0.02	0.75	0.08
	10<E	0.04	0.04	2.03	0.11
Neutrons	E<10	0.0002	0.0003	0.0023	0.0003
	10<E	0.0047	0.0098	0.0039	0.0012

Type	E range (MeV)	Ca48 @ 80 uA	LD2 @ 50 uA	PREX_I @ 70 uA	PREX_II @ 70 uA
		Power (W)	Power (W)	Power (W)	Power (W)
e±	E<10	9.02	5.67	11.64	0.64
	10<E	14.45	8.13	90.16	3.93
Photons	E<10	1.94	1.13	52.19	5.34
	10<E	3.11	2.05	142.31	7.49
Neutrons	E<10	0.02	0.02	0.16	0.02
	10<E	0.38	0.49	0.28	0.08

Radiation Flux with No Poly

Type	E range (MeV)	Ca48	LD2	PREX I	PREX II
		Flux (Hz/uA)	Flux (Hz/uA)	Flux (Hz/uA)	Flux (Hz/uA)
e±	E<10	3.83E+11	4.08E+11	3.45E+11	2.05E+10
	10<E	3.84E+10	3.23E+10	2.10E+11	9.42E+09
Photons	E<10	2.22E+11	2.83E+11	5.49E+12	6.41E+11
	10<E	6.77E+09	6.70E+09	3.43E+11	1.93E+10
Neutrons	E<10	7.86E+08	1.44E+09	1.62E+10	1.79E+09
	10<E	2.62E+08	3.31E+08	3.93E+08	1.34E+08

Type	E range (MeV)	Ca48 @ 80 uA	LD2 @ 50 uA	PREX_I @ 70 uA	PREX_II @ 70 uA
		Flux (Hz)	Flux (Hz)	Flux (Hz)	Flux (Hz)
e±	E<10	3.06E+13	2.04E+13	2.42E+13	1.44E+12
	10<E	3.07E+12	1.62E+12	1.47E+13	6.59E+11
Photons	E<10	1.78E+13	1.41E+13	3.85E+14	4.49E+13
	10<E	5.41E+11	3.35E+11	2.40E+13	1.35E+12
Neutrons	E<10	6.29E+10	7.21E+10	1.13E+12	1.25E+11
	10<E	2.10E+10	1.65E+10	2.75E+10	9.39E+09

09/09/15

Rakitha Beminiwattha

3

- Nuclear modification has many open important questions for our understanding of QCD and can be tested with SoLID PVDIS
- Working on improving proposal to address PAC concerns
- Updates need to be made on target and radiation estimates

BACKUP

$$R^{\gamma(\gamma Z)} \equiv \frac{\sigma_L^{\gamma(\gamma Z)}}{\sigma_T^{\gamma(\gamma Z)}} = r^2 \frac{F_2^{\gamma(\gamma Z)}}{F_1^{\gamma(\gamma Z)}} - 1$$

$$r^2 = 1 + \frac{Q^2}{\nu} = 1 + \frac{4M^2 x^2}{Q^2}$$

$$A_{PV} = - \left(\frac{G_F Q^2}{4\sqrt{2}\pi\alpha} \right) \left[g_A^e Y_1 \frac{F_1^{\gamma Z}}{F_1^\gamma} + \frac{g_V^e}{2} Y_3 \frac{F_3^{\gamma Z}}{F_1^\gamma} \right]$$

$$Y_1 = \frac{1 + (1-y)^2 - y^2 (1 - r^2/(1 + R^{\gamma Z})) - 2xyM/E}{1 + (1-y)^2 - y^2 (1 - r^2/(1 + R^\gamma)) - 2xyM/E} \left(\frac{1 + R^{\gamma Z}}{1 + R^\gamma} \right)$$

$$Y_3 = \frac{1 - (1-y)^2}{1 + (1-y)^2 - y^2 (1 - r^2/(1 + R^\gamma)) - 2xyM/E} \left(\frac{r^2}{1 + R^\gamma} \right)$$

$$F_1^\gamma = \frac{1}{2} \sum_i e_i^2 (q_i(x) + \bar{q}_i(x)); F_2^\gamma = 2xF_1^\gamma,$$

$$F_1^{\gamma Z} = \sum_i e_i g_V^i (q_i(x) + \bar{q}_i(x)); F_2^{\gamma Z} = 2xF_1^{\gamma Z},$$

$$F_3^{\gamma Z} = 2 \sum_i e_i g_A^i (q_i(x) - \bar{q}_i(x)).$$

Possible future extensions


Experiment	Constraint	Reason
LD ₂	CSV, Q^2 dependence	
LH ₂	d/u	
⁴⁸ Ca	Isovector EMC properties	
⁴⁰ Ca	$R^{\gamma Z}$, CSV dependence on A	Effects are very large or in serious disagreement with NuTeV or other data
Co, Cu?	Less direct test on NuTeV	Effects are very large or in serious disagreement with NuTeV or other data
⁵⁶ Fe	Direct test of NuTeV	Separate data point
²⁰⁸ Pb	Very high Z	Separate data point

Approved

This Proposal


Can't with SoLID


- Varying weights in fits between lepton/Drell Yan and ν can show tension between data sets
- nCTEQ fits show dramatic differences in a similar vein at CBT
- Few percent effect in asymmetry




- w relative weighting of ν data with lepton/DY
- Suggests softness in this channel through lepton/DY
- PVDIS can provide differentiation

- Nuclear dependence of $R^{\gamma Z}$ data doesn't exist





π^- to e^- Ratio for ^{48}Ca and LD Target



GEM plane	LD ₂ background (kHz/mm ² /μA)	⁴⁸ Ca EM background (kHz/mm ² /μA)	⁴⁸ Ca EM background (no baffles) (kHz/mm ² /μA)
1	6.8	4.8	49.4
2	3.0	2.1	32.3
3	1.1	0.8	9.9
4	0.7	0.5	6.4

Momentum range (GeV)	π^- (MHz)	π^+ (MHz)	$\pi^0(\gamma)$ (MHz)	Proton (MHz)	EM (γ, e^\pm) (GHz)
p > 0.0 GeV	618	283	70123	483	844
p > 0.3 GeV	439	153	438	417	n/a
p > 1.0 GeV	123	18	37	51	0.0
p > 3.0 GeV	2	0.01	0.04	0.004	0.0

ECal Trigger Rates

region	full	high	low
rate entering the EC (kHz)			
e^-	240	129	111
π^-	5.9×10^5	3.0×10^5	3.0×10^5
π^+	2.7×10^5	1.5×10^5	1.2×10^5
$\gamma(\pi^0)$	7.0×10^7	3.5×10^7	3.5×10^7
p^+	4.8×10^5	2.1×10^5	2.7×10^5
sum	7.1×10^7	3.6×10^7	3.6×10^7
Rate for $p < 1$ GeV (kHz)			
sum	8.4×10^8	4.2×10^8	4.2×10^7
trigger rate for $p > 1$ GeV (kHz)			
e^-	152	82	70
π^-	4.0×10^3	2.2×10^3	1.8×10^3
π^+	0.2×10^3	0.1×10^3	0.1×10^3
$\gamma(\pi^0)$	3	3	0
p	1.6×10^3	0.9×10^3	0.7×10^3
sum	5.9×10^3	3.3×10^3	2.6×10^3
trigger rate for $p < 1$ GeV (kHz)			
sum	2.8×10^3	1.4×10^3	1.4×10^3
Total trigger rate (kHz)			
total	8.7×10^3	4.7×10^3	4.0×10^3

Cerenkov Trigger Rates


	Total Rate for $p > 0.0$ GeV (kHz)	Rate for $p > 3.0$ GeV (kHz)
DIS	240	73
π^-	5.9×10^5	1.6×10^3
π^+	2.7×10^5	40
$\gamma(\pi^0)$	7.0×10^7	40
ρ	4.8×10^5	4
Sum	7.1×10^7	1.7×10^3
Trigger Rate from Cherenkov (kHz)		
	Trigger Rate for $p > 1.0$ GeV (kHz)	Trigger Rate for $p > 3.0$ GeV (kHz)
DIS	223	66
π^-	193	49
π^+	22	1.6
$\gamma(\pi^0)$	0	0
ρ	0	0
Sum	438	116

Radiation Type	E-Range (MeV)	Incident Radiation Power	
		^{48}Ca (W/ μA)	LD ₂ (W/ μA)
e^{\pm}	E < 10	0.13	0.13
	E > 10	0.19	0.17
n	E < 10	0.0001	0.0006
	E > 10	0.02	0.04
γ	E < 10	0.02	0.02
	E > 10	0.04	0.05

Rates and Backgrounds


- Trigger defined by coincidence between Cherenkov and shower - 150 kHz total anticipated with background (well below SoLID spec)
- Pion contamination no worse than 4% in any given bin (worst at high x)
- GEM rates comparable to or smaller than design for LD₂

EM Background Rate in the GEM Detectors



Particle	DAQ Coin.	Trig. Rate (kHz)
	P > 1 GeV	P > 3 GeV
DIS e^-	144	61
π^-	11	7
π^+	0.4	0.2
Total	155	68

- Many potential nuclear effects come into play as this sector is not presently well constrained
- Requires measurements from LD₂ and LH₂ for information on size of nuclear effects
- Free PDFs will be constrained by LH₂ et al



- Many potential nuclear effects come into play as this sector is not presently well constrained
- Requires measurements from LD₂ and LH₂ for information on size of nuclear effects
- Higher twist effects will also be constrained by LD₂ using same kinematics, but also 6.6 GeV beam

Approved LD₂ (120 days):

