

## Single-Transverse-Spin Asymmetries of Identified Charged Hadrons in Polarized $pp$ Collisions at $\sqrt{s} = 62.4$ GeV

I. Arsene,<sup>9</sup> I. G. Bearden,<sup>4</sup> D. Beavis,<sup>1</sup> S. Bekele,<sup>8</sup> C. Besliu,<sup>7</sup> B. Budick,<sup>3</sup> H. Bøggild,<sup>4</sup> C. Chasman,<sup>1</sup> H. H. Dalsgaard,<sup>4</sup> R. Debbé,<sup>1</sup> B. Fox,<sup>1</sup> J. J. Gaardhøje,<sup>4</sup> K. Hagel,<sup>5</sup> A. Jipa,<sup>7</sup> E. B. Johnson,<sup>8</sup> R. Karabowicz,<sup>2</sup> N. Katryńska,<sup>2</sup> E. J. Kim,<sup>8</sup> T. M. Larsen,<sup>4</sup> J. H. Lee,<sup>1</sup> G. Løvhøiden,<sup>9</sup> Z. Majka,<sup>2</sup> M. Murray,<sup>8</sup> C. Nygaard,<sup>4</sup> J. Natowitz,<sup>5</sup> B. S. Nielsen,<sup>4</sup> D. Pal,<sup>8</sup> A. Qviler,<sup>9</sup> C. Ristea,<sup>4</sup> D. Röhrich,<sup>6</sup> S. J. Sanders,<sup>8</sup> P. Staszczel,<sup>2</sup> T. S. Tveter,<sup>9</sup> F. Videbæk,<sup>1</sup> H. Yang,<sup>6</sup> and R. Wada<sup>5</sup>

(BRAHMS Collaboration)

<sup>1</sup>Brookhaven National Laboratory, Upton, New York 11973, USA

<sup>2</sup>Smoluchowski Inst. of Physics, Jagiellonian University, Krakow, Poland

<sup>3</sup>New York University, New York, New York 10003, USA

<sup>4</sup>Niels Bohr Institute, Blegdamsvej 17, University of Copenhagen, Copenhagen 2100, Denmark

<sup>5</sup>Texas A&M University, College Station, Texas 77843, USA

<sup>6</sup>Department of Physics, University of Bergen, Bergen, Norway

<sup>7</sup>University of Bucharest, Bucharest, Romania

<sup>8</sup>University of Kansas, Lawrence, Kansas 66049, USA

<sup>9</sup>Department of Physics, University of Oslo, Oslo, Norway

(Received 7 January 2008; published 22 July 2008)

The first measurements of  $x_F$ -dependent single-spin asymmetries of identified charged hadrons,  $\pi^\pm$ ,  $K^\pm$ , and protons, from transversely polarized proton-proton collisions at 62.4 GeV at RHIC are presented. Large asymmetries are seen in the pion and kaon channels. The asymmetries in inclusive  $\pi^+$  production,  $A_N(\pi^+)$ , increase with  $x_F$  from 0 to  $\sim 0.25$  and  $A_N(\pi^-)$  decrease from 0 to  $\sim -0.4$ . Observed asymmetries for  $K^-$  unexpectedly show positive values similar to those for  $K^+$ , increasing with  $x_F$ , whereas proton asymmetries are consistent with zero over the measured kinematic range. Comparisons of the data with predictions of QCD-based models are presented.

DOI: 10.1103/PhysRevLett.101.042001

PACS numbers: 13.85.Ni, 12.38.Qk, 13.88.+e

The transverse spin dependence of hadron cross sections in  $p^\uparrow + p$  ( $\bar{p}^\uparrow + p$ ) reactions in the energy regime where perturbative QCD (pQCD) is applicable are expected to be negligibly small [1] in the lowest-order QCD approximation, whereas experimentally large left-right asymmetries have been observed [2–4] for large Feynman- $x$ ,  $x_F = 2p_L/\sqrt{s}$ . Measurements of large asymmetries of inclusive pion production and polarization in the production of hyperons [5] in a wide energy range have motivated various theoretical efforts to understand the phenomena. Observed large asymmetries are not particularly new phenomena in  $p^\uparrow + p$  reactions since sizable asymmetries in inclusive pion production had been observed in the lower energy regime [6,7], but understanding asymmetries in hadron reactions where partonic QCD descriptions are more relevant poses a new theoretical challenge. The main theoretical focus to account for the observed single-spin asymmetries (SSAs) in the framework of QCD has been on the role of transverse momentum dependent (TMD) partonic effects in the structure of the initial transversely polarized nucleon (“Sivers” mechanism) [8] and on the fragmentation process of a polarized quark into hadrons (“Collins” mechanism) [9]. Higher twist effects (“twist-3”) arising from quark-gluon correlation effects beyond the conventional twist-2 distribution have also

been considered as a possible origin of SSA [10,11]. SSA measurements in  $p^\uparrow + p$  at RHIC energies are of particular interest because the next-to-leading-order (NLO) pQCD calculations [12] for the unpolarized (spin-averaged) meson cross sections at forward rapidities successfully describe the data [13,14]. At  $\sqrt{s} \sim 20$  GeV where Fermi National Accelerator Laboratory–E704 observed large SSAs, NLO pQCD cross-section calculations [12] significantly underpredict the measurements showing increasing discrepancies with increasing  $x_F$  [15]. The disagreements indicate that there may be another mechanism, likely related to “soft” processes, which is significantly responsible for pion production at this lower energy. The two sets of data at  $\sqrt{s} \sim 20$  GeV and at  $\sqrt{s} = 200$  GeV cover a similar kinematic range in  $x_F$  and  $p_T$  and the measurements show that SSAs for pions are energy independent to first approximation. Since pQCD description of cross sections at the high-energy region is quite successful, while it fails in the low energy domain, it might imply that the dominant mechanism responsible for the large SSAs at the two different energies is a manifestation of two different phenomena. The newly available measurements from RHIC in the intermediate energy regime at  $\sqrt{s} = 62.4$  GeV in  $p^\uparrow + p$  can uniquely provide an opportunity to clarify the pQCD contribution to SSAs and their energy

dependences. A simultaneous description of SSAs and the unpolarized cross sections [16] in a wide kinematic range will be a crucial test for the partonic pQCD description. In particular, flavor dependent SSA measurements allow more complete and stringent tests of theoretical models due to flavor dependence in parton distribution functions and fragmentation processes. We present here the first measurement of  $x_F$ -dependent SSAs of identified charged hadrons,  $\pi^\pm$ ,  $K^\pm$ , and protons, from transversely polarized proton-proton collisions at 62.4 GeV at RHIC.

The SSA is defined as a “left-right” asymmetry of produced particles from the hadronic scattering of transversely polarized protons by unpolarized protons. Experimentally the asymmetry can be obtained by flipping the spins of polarized protons, and is customarily defined as analyzing power  $A_N$ :

$$A_N = \frac{1}{\mathcal{P}} \frac{(N^+ - \mathcal{L}N^-)}{(N^+ + \mathcal{L}N^-)}, \quad (1)$$

where  $\mathcal{P}$  is polarization of the beam,  $\mathcal{L}$  is the spin dependent relative luminosity ( $\mathcal{L} = \mathcal{L}_+/\mathcal{L}_-$ ) and  $N^{+(-)}$  is the number of detected particles with beam spin vector oriented up (down). Since both colliding beams are polarized at RHIC, the polarization of “target” protons is averaged over in Eq. (1). The systematic error on the  $A_N$  measurements is estimated to be 10% including uncertainties from the beam polarization,  $\delta\mathcal{P}/\mathcal{P} \sim 7.2\%$  for the “blue” beam (circulating clockwise) and 9.3% for the “yellow” beam (circulating counterclockwise). The polarization of the blue (yellow) beam is utilized for the  $A_N$  measurements of particles in positive (negative)  $x_F$ . The systematic error represents mainly scaling uncertainties on the values of  $A_N$ . The average polarization of the beam  $\mathcal{P}$  measured by the hydrogen jet and pC polarimeters is about 50% for the blue and yellow beams [17].

The data presented here were collected by the BRAHMS detector system [18] with polarized  $p + p$  collisions from RHIC with a sampled integrated luminosity of  $0.21 \text{ pb}^{-1}$  at  $\sqrt{s} = 62.4 \text{ GeV}$ . The relative luminosity ( $\mathcal{L}$ ) between the sums of spin-up and spin-down bunches was measured with a set of Cherenkov radiators placed symmetrically with respect to the nominal interaction point [14]. The detectors cover the pseudorapidity ( $\eta$ ) interval from  $3.26 < |\eta| < 5.25$ , and are measured from Vernier scans to be sensitive to  $\sim 33\%$  of the total inelastic cross section of 36 mb at 62.4 GeV. The uncertainty of determining the relative luminosities is estimated to be 0.3%. The forward spectrometer (FS) measures charged particle tracks in the forward kinematic region ( $\theta = 2.3^\circ - 15^\circ$ ) with good momentum resolution and particle identification. The momentum ( $p$ ) resolution of the FS is  $\delta p/p \approx 0.0016p$  for the half field setting where  $p$  is in GeV/ $c$ . Particle identification was done by utilizing the Ring Image Cherenkov Detector (RICH) [19] detector which is capable of identifying pions and kaons up to  $p \sim 35 \text{ GeV}/c$  and protons

above  $17 \text{ GeV}/c$  with an efficiency of  $\sim 97\%$  and a negligible ( $\leq 0.5\%$ ) probability of misidentification in the measured kinematic range ( $p < 20 \text{ GeV}/c$ ). The kinematic coverages of the data taken with the FS at  $2.3^\circ$ ,  $3^\circ$ , and  $6^\circ$  as a function of  $p_T$  and  $x_F$  are shown in Fig. 1, where the narrow  $p_T - x_F$  correlated band at a given setting is due to the small aperture of the spectrometer. A detailed description of the spectrometer and other experimental details can be found in [18].

The analyzing power  $A_N$  for charged pions,  $A_N(\pi^+)$  and  $A_N(\pi^-)$  at  $\sqrt{s} = 62.4 \text{ GeV}$  as a function of  $x_F$  is shown in Fig. 2 for the two FS angle settings,  $2.3^\circ$  and  $3^\circ$ . At a fixed  $x_F$  value, the  $3^\circ$  setting samples higher  $p_T$  pions as indicated in Fig. 1. The mean  $p_T$  values  $\langle p_T \rangle$  at  $x_F = 0.55$  are 1.08 and 1.28 GeV/ $c$  at  $2.3^\circ$  and at  $3^\circ$ , respectively [20]. The measured  $A_N$  values show strong dependence in  $x_F$  reaching large asymmetries up to  $\sim 40\%$  at  $x_F \sim 0.6$  and no significant asymmetries at  $-x_F$ . The decrease of  $A_N$  at high- $p_T$  ( $\geq 1 \text{ GeV}/c$ ) and high- $x_F$ , especially for  $\pi^+$ , as shown in Fig. 2 by comparing the two sets of measurements at  $2.3^\circ$  and at  $3^\circ$  might indicate that  $A_N$  is in accordance with the expected power-suppressed nature of  $A_N$  [21]. The asymmetries and their  $x_F$  dependence are qualitatively in agreement with the measurements from E704 at  $\sqrt{s} = 19.3 \text{ GeV}$  and also most recent  $A_N(\pi^0)$  measurements at RHIC  $\sqrt{s} = 200 \text{ GeV}$  [3,13]. Figure 2 also compares  $A_N(\pi)$  with a pQCD calculation in the range of  $p_T > 1 \text{ GeV}/c$  using “extended” twist-3 parton distributions [10] including the “nonderivative” contributions [21–23]. In this framework, results of two calculations from the model are compared with the data. One is with only two quark valence densities ( $u_v, d_v$ ) in the ansatz, which is shown in Fig. 2. The second with additional sea- and anti- quark contributions in the model fit slightly




FIG. 1 (color online).  $p_T$  vs  $x_F$  for the data used in the SSA analysis. The dotted boxes are for the measurements from FS at  $6^\circ$ , the filled boxes are from FS at  $2.3^\circ$  and the empty boxes with solid line are from  $3^\circ$ . Data from FS at  $2.3^\circ$  and  $3^\circ$  are used in combination for kaons and protons. The size of the boxes represents the relative intensity of the data in logarithmic scale. The 5 bands marked as (a)–(e) are the  $p_T$  ranges used in Fig. 3.




FIG. 2 (color online).  $A_N$  vs  $x_F$  for  $\pi^+$  and  $\pi^-$ . Circle symbols are for  $\pi^+$  and box symbols are for  $\pi^-$  measured in FS at  $2.3^\circ$  (solid symbols) and  $3^\circ$  (open symbols). The curves are from theoretical calculations. Solid lines are to be compared with the data at  $2.3^\circ$  and dotted lines are for  $3^\circ$ . Thick (solid and dotted) lines are from the initial-state twist-3 calculations [21,23], medium lines are from the final-state twist-3 calculations [24,34]. Predictions from the Siverson function calculations are shown as thin lines [25,26]. Only statistical errors are shown where larger than symbols.

increases  $A_N(\pi)$  ( $\sim 5\%$ ). As the calculations show, the dominant contribution to SSA is from valence quarks with contributions from sea- and anti- quarks small enough that the current measurements are not able to quantitatively constrain the contribution. The calculations, which were done in the same kinematic range as the data, describe the data, especially  $A_N(\pi^-)$  within the uncertainties.  $A_N(\pi)$  calculated from the “final-state twist-3” [24] which uses the twist-3 fragmentation function (FF) for the pion clearly under-predicts  $A_N(\pi^-)$  while is in a reasonable agreement within uncertainties for  $A_N(\pi^+)$ . In Fig. 2, the data are also compared with calculations including Siverson mechanism which successfully describe the E704  $A_N$  data using valence-like Siverson functions [25,26] for  $u$  and  $d$  quarks with opposite sign. The FFs used are from the KKP parameterization [27], but the Kretzer FF [28] gives similar results. The calculations underestimate  $A_N$ , which indicates that TMD parton distributions are not sufficient to describe the SSA data at this energy. As very recent studies [29] suggest, Collins mechanism might also be needed to account

fully for the observed asymmetries. All  $A_N(\pi)$  calculations compared with the data shows  $|A_N(\pi^+)| \sim |A_N(\pi^-)|$  while the data exhibit  $|A_N(\pi^+)| < |A_N(\pi^-)|$  where  $p_T \gtrsim 1$  GeV/c. Since there is a strong kinematic correlation between  $x_F$  and  $p_T$  in the data as shown in Fig. 1, the rise of  $A_N$  in Fig. 2 can be also driven by  $p_T$ .

Figure 3 shows  $A_N(\pi^+)$  and  $A_N(\pi^-)$  for 5 different  $p_T$  regions from 0.4 to 1.2 GeV/c. As seen in Fig. 3, the  $x_F$  dependence of  $A_N$  at low- $p_T$  ( $p_T \lesssim 0.5$  GeV/c) is very small but increases with  $p_T$  in the kinematic region at least up to  $p_T \sim 1$  GeV/c. The  $p_T$  dependence of analyzing powers with  $x_F$  is qualitatively consistent with the measurements at  $\sqrt{s} = 19.3$  GeV, where strong  $x_F$  dependent SSAs is observed only above a  $p_T$  “threshold” ( $\lesssim 0.7$  GeV/c) [3]. It is noted that the trend is also qualitatively in agreement with the polarization of the  $\Lambda_s$  produced at the same collision energy,  $\sqrt{s} = 62$  GeV [5].

The SSAs for charged kaons as a function of  $x_F$  are shown in Fig. 4 together with twist-3 and Siverson calculations (see the figure caption for details). The asymmetry for  $K^+(u\bar{s})$  is positive as is the  $A_N$  of  $\pi^+(ud)$ , which is expected if the asymmetry is mainly carried by valence quarks, but the measured positive SSAs of  $K^-(\bar{u}s)$  seem to contradict the naïve expectations [30] of valence quark dominance. In a valence-like model (no Siverson effect from sea-quarks and/or gluons), nonzero positive  $A_N(K^-)$  implies large nonleading FFs ( $D_u^{K^-}$ ,  $D_d^{K^-}$ ) and insignificant contribution from strange quarks. Twist-3 calculations using Kretzer FF also under-predict  $A_N(K^-)$  due to the small contribution of sea and strange-quark contribution to  $A_N$  in the model. Notably the unpolarized cross section for  $K^+$  is an order of magnitude higher than for  $K^-$  [16]. The current calculations for kaon asymmetries need an extra or a different mechanism to account for positive  $A_N(K^-)$  at similar level of  $A_N(K^+)$  as shown in Fig. 4. If the asymmetries of  $K^-$  are mainly driven by pQCD effects, the discrepancies between data and calculations are expected to be reduced once the Siverson function is better understood for sea quarks and also FFs especially the unfavored FFs. Likewise possible non-negligible contributions from the Collins mechanism, as recently reported [31,32], may need to be explored further.




FIG. 3 (color online).  $A_N$  vs  $x_F$  for  $\pi^+$  and  $\pi^-$  for positive  $x_F$  at fixed  $p_T$  values: (a)  $0.4 < p_T < 0.5$ , (b)  $0.5 < p_T < 0.6$ , (c)  $0.6 < p_T < 0.8$ , (d)  $0.8 < p_T < 1.0$ , and (e)  $1.0 < p_T < 1.2$  GeV/c as shown in Fig. 1, respectively.




FIG. 4 (color online).  $A_N(K^+)$  and  $A_N(K^-)$  vs  $x_F$ . Circle symbols are for  $K^+$  and box symbols are for  $K^-$ . The solid ( $K^+$ ) and dotted ( $K^-$ ) lines are from the initial-state twist-3 calculations with (thick lines) and without (medium lines) sea- and antiquark contribution. Calculations for the Sivers function are shown as thin lines. Errors are statistical only.

SSAs at  $x_F < 0$  probe the kinematics of the sea (gluon) region of  $p^\uparrow$  at small- $x$  and the valence region of  $p$ , which was experimentally measured by the produced particles in the forward hemisphere of  $p$  in the  $p + p^\uparrow$  collisions utilizing the polarization information of the target. The measured insignificant  $A_N$  for pions and kaons in large  $|x_F|$  when  $x_F < 0$  indicates no significant contribution to  $A_N$  from processes where  $gq$  scattering is enhanced, and the asymmetries are dominated by the processes where large quark PDFs and FFs are expected. In Fig. 5, we demonstrate that inclusive protons show no significant asymmetries in contrast to pions and kaons in the forward kinematic region. The insignificant asymmetries observed are consistent with the measurements at lower energies [2,33], but require more understanding of their production mechanism to theoretically describe the behavior because a significant fraction of the protons might still be related to the polarized beam fragments at this kinematic range [14].

In summary, BRAHMS has measured SSAs for inclusive identified charged hadron production at forward rapidities in  $p^\uparrow + p$  at  $\sqrt{s} = 62.4$  GeV. A twist-3 pQCD model describes the  $x_F$  dependence of  $A_N(\pi)$  and the energy dependence at high- $p_T$  ( $p_T > 1$  GeV/ $c$ ) where the calculations are applicable, but it remains a challenge for pQCD models to consistently describe spin-averaged cross sections at this energy [15,16]. Measurements of  $A_N$




FIG. 5 (color online).  $A_N$  vs  $x_F$  of the proton.

for kaons and protons suggest the possible manifestation of non-pQCD phenomena and call for more theoretical modeling with improved understanding of the fragmentation processes. The energy and flavor dependent asymmetry measurements impose an important constraint on theoretical models describing fundamental mechanisms of transverse spin asymmetries and the quantum chromodynamical description of hadronic structure.

We thank F. Yuan, U. D'Alesio and Y. Koike for providing us with their calculations. This work was supported by the office of NP in DoE (USA), NSRC (Denmark), RC (Norway), SCSR (Poland), MoR (Romania), and a sponsored research grant from Renaissance Technologies Corp.

- [1] G.L. Kane *et al.*, Phys. Rev. Lett. **41**, 1689 (1978).
- [2] S. Saroff *et al.*, Phys. Rev. Lett. **64**, 995 (1990).
- [3] D.L. Adams *et al.*, Phys. Lett. B **264**, 462 (1991).
- [4] A. Bravar *et al.*, Phys. Rev. Lett. **77**, 2626 (1996).
- [5] S.M. Smith *et al.*, Phys. Lett. B **185**, 209 (1987).
- [6] W.H. Dragoset, Jr. *et al.*, Phys. Rev. D **18**, 3939 (1978).
- [7] J. Antille *et al.*, Phys. Lett. B **94**, 523 (1980).
- [8] D. Sivers, Phys. Rev. D **41**, 83 (1990).
- [9] J.C. Collins, Nucl. Phys. **B396**, 161 (1993).
- [10] J. Qiu and G. Sterman, Phys. Rev. D **59**, 014004 (1998).
- [11] J. Kodaira and K. Tanaka, Prog. Theor. Phys. **101**, 191 (1999).
- [12] B. Jager *et al.*, Phys. Rev. D **67**, 054005 (2003).
- [13] J. Adams *et al.*, Phys. Rev. Lett. **92**, 171801 (2004).
- [14] I. Arsene *et al.*, Phys. Rev. Lett. **98**, 252001 (2007).
- [15] C. Bourrely and J. Soffer, Eur. Phys. J. C **36**, 371 (2004).
- [16] BRAHMS Collaboration, Cross-section measurements of identified hadrons at  $\sqrt{s} = 62.4$  GeV (to be published).
- [17] A. Bazilevsky *et al.*, RHIC/CAD/AP Note 298 (2007).
- [18] M. Adamczyk *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **499**, 437 (2003).
- [19] R. Debbé *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **570**, 216 (2007).
- [20] The data are available in a tabulated form in <http://www.rhic.bnl.gov/brahms/WWW/publications.html>.
- [21] C. Kouvaris *et al.*, Phys. Rev. D **74**, 114013 (2006).
- [22] W. Vogelsang, Proc. of 17th International Spin Physics Symposium, SPIN2006, Oct. 2006, Kotyo, Japan AIP Conf Proc. No. 915.
- [23] The calculations were provided by Feng Yuan.
- [24] Y. Koike, Proc. of 15th International Spin Physics Symposium, SPIN2002, Sept. 2002, Upton, USA AIP Conf Proc. No. 675.
- [25] U. D'Alesio and F. Murgia, Phys. Rev. D **70**, 074009 (2004).
- [26] The calculations were provided by Umberto D'Alesio.
- [27] B.A. Kniehl *et al.*, Nucl. Phys. **B597**, 337 (2001).
- [28] S. Kretzer, Phys. Rev. D **62**, 054001 (2000).
- [29] F. Yuan, arXiv:0801.3441; arXiv:0804.3047.
- [30] M. Anselmino *et al.*, Phys. Lett. B **442**, 470 (1998).
- [31] E. S. Ageev *et al.*, Nucl. Phys. **B765**, 31 (2007).
- [32] M. Diefenthaler, arXiv:0706.2242.
- [33] V. Abramov *et al.*, Proc. of 16th International Spin Physics Symposium, SPIN2004, Oct. 2004, Trieste, Italy (World Scientific, Singapore, 2005).
- [34] The calculations were provided by Yuji Koike.