Precision Measurement of Neutron Asymmetry A_1^n at Large x_{Bj}

X. Zheng, M.I.T., Cambridge, MA

BNL Spin Discussion, May 21, Upton, NY

Outline

- Introduction - Polarized DIS and Spin Structure Functions
- Theories about A_1^n
- Experiment E99-117
  - Experimental Setup
  - Jefferson Lab Hall A Polarized ^{3}He Target
  - Data Analysis – From ^{3}He to Neutron
- Preliminary Results
- Discussion, Summary and Outlook

List of Collaborators

J. Gao
California Institute of Technology
K. Aniol, D. Margaziotis
California State University, LA
P. Markowitz
Florida International University
R. Roche
Florida State University
M. Roedelbronn
University of Illinois
F. Cusanno, R. De Leo, F. Garibaldi, S. Frullani, G. Urciuoli
INFN
J.-P. Chen, E. Chudakov, J. Gomez, K. de Jager, R. Michaels, O. Hansen,
J. LeRose, N. Liyanage, B. Reitz, A. Saha, B. Wojtsekhowski
Jefferson Lab
K. McCormick
Kent State University
W. Korsch, P. Zolnierczuk
University of Kentucky
J. Kelly, T. Horn, N. Savvinov
University of Maryland
L. Kaufman, A. Vacheret
University of Massachussetts
W. Bertozzi, Z. Chai, D. Dutta, H. Gao, D. Higinbotham, M. Rvachev, S. Sirca,
H. Xiang, Y. Xiao, F. Xiong, B. Zhang, X. Zheng, L. Zhu
Massachussetts Institute of Technology
J. Calarco
University of New Hampshire
W. Hinton
Old Dominion University
E. Busato, S. Dieterich, R. Gilman, X. Jiang, S. Strauch
Rutgers University
P. Souder
Syracuse University
G. Ron
Tel Aviv
Z.-E. Meziani, S. Choi, K. Slifer, P. Solvignon
Temple University
A. Camsonne, G. Cates, A. Deur, J. Singh
University of Virginia
D. Armstrong, T. Averett, K. Kramer, S. Binet, C. Butuceanu, M. Finn,
B. Moffit, S. Phillips, A. Powell, J. Roche, D. Steiner, V. Sulkosky, X. Zhu
College of William and Mary

Polarized Deep Inelastic Scattering

POLARIZED DIS



STRUCTURE FUNCTIONS

- Unpolarized Target:

$$\frac{d^2\sigma_{\uparrow\Uparrow}}{d\Omega dE'} + \frac{d^2\sigma_{\uparrow\Downarrow}}{d\Omega dE'} = \frac{8\alpha^2 E'^2}{mQ^4}[2\sin^2\frac{\theta}{2}F_1(x,Q^2) + \frac{m^2}{\nu}\cos^2\frac{\theta}{2}F_2(x,Q^2)]$$

$$F_1(x,Q^2) = \frac{F_2(x,Q^2)(1+\gamma^2)}{2x(1+R(x,Q^2)}$$

- Longitudinally Polarized Target:

$$\frac{d^2\sigma_{\uparrow\Uparrow}}{d\Omega dE'} - \frac{d^2\sigma_{\uparrow\Downarrow}}{d\Omega dE'} = \frac{4\alpha^2 E'}{E\nu Q^2}\Big[(E+E'\cos\theta)g_1(x,Q^2) - 2xmg_2(x,Q^2)\Big]$$

- Transversely Polarized Target:

$$\frac{d^2\sigma_{\uparrow\Rightarrow}}{d\Omega dE'} - \frac{d^2\sigma_{\uparrow\Leftarrow}}{d\Omega dE'} = \frac{4\alpha^2 E'^2}{E\nu Q^2}\sin\theta\Big[g_1(x,Q^2) + \frac{2Em}{\nu}g_2(x,Q^2)\Big]$$

Structure Functions in QPM

Within the Quark-Parton Model,

$$F_1(x) = \frac{1}{2}\sum_i e_i^2\left[q_i^{\uparrow}(x) + q_i^{\downarrow}(x)\right]$$

$$g_1(x) = \frac{1}{2}\sum_i e_i^2\left[q_i^{\uparrow}(x) - q_i^{\downarrow}(x)\right]$$

Ignoring quark mass effects,

$$g_2(x,Q^2) = g_2^{WW}(x,Q^2) + \bar{g}_2(x,Q^2)$$

$$g_2^{WW}(x,Q^2) = -g_1(x,Q^2) + \int_x^1 \frac{g_1(y,Q^2)}{y}\mathrm{dy}$$

Virtual Photon Asymmetries

$$A_1 = \frac{\sigma_{\frac{1}{2}} - \sigma_{\frac{3}{2}}}{\sigma_{\frac{1}{2}} + \sigma_{\frac{3}{2}}}$$



$$A_1(x,Q^2) = \frac{g_1(x,Q^2) - \gamma^2 g_2(x,Q^2)}{F_1(x,Q^2)}$$

A_1^n at high x_{Bj}

WHY AT HIGH x_{Bj}?

- Valence quarks dominate;
- Constituent quark models
  - Is our understanding of valence quark picture correct?
- pQCD
  - how solid it is and to how low x it can apply;



Theories about A_1^n

- SU(6): $A_1^n = 0$
- SU(6) Broken:$A_1^n \to 1$ as $x \to 1$;

- Constituent quark models (CQM): N.Isgur $A_1^n \to 1$ as $x \to 1$;
- pQCD at $Q^2 \to \infty$, $A_1^n \to 1$ as $x \to 1$;

- Statistical model, C.Bourrely, J. Soffer ;
- Duality: W. Melnitchouk;

- Chiral Quark-Soliton Model, H. Weigel, $A_1^n < 0$ for $x > 0.5$;
- Instanton Model, N.I.Kochelev, $A_1^n \sim 0$ or $A_1^n < 0$;

- CQM: N.Isgur, Phys.Rev. **D59** (1999) 034013
  - Hyperfine perturbed
  - A (simple) parameterization: $\frac{d(x)}{u(x)} \approx \kappa(1-x)$ as $x \to 1,\ \kappa \approx 0.5 \sim 0.6$
    $c_A(x) = nx(1-x)^n,\ n = 2 \sim 4$
  - $A_1^n(x) = \frac{4\Delta d_v(x)+\Delta u_v(x)}{4d_v(x)+u_v(x)}$ $A_1^p(x) = \frac{4\Delta u_v(x)+\Delta d_v(x)}{4u_v(x)+d_v(x)}$
  - A_1^n and $A_1^p \to 1$ as $x \to 1$.

World data on A_1^n and prediction from CQM



- pQCD (LSS-BBS): E. Leader, A.V.Sidorov, D.B.Stamenov, hep-ph/9708335
  - Polarized & unpolarized parton densities by Brodsky, Burkhardt and Schmidt (BBS);
  - $\frac{\Delta u}{u} \to 1$, $\frac{\Delta d}{d} \to 1$ at $x \to 1$.
- pQCD (LSS2001): E. Leader, A.V.Sidorov, D.B.Stamenov, hep-ph/0111267
  - A new evaluation of NLO polarized parton densities in the nucleon;
  - calculation done for g_1/F_1 at Q^2= 3, 5, 100 (GeV/c)^2.

World data on A_1^n and pQCD predictions



- Statistical Model: C.Bourrely, J. Soffer, hep-ph/0109160
  - global NLO QCD analysis of unpolarized and polarized DIS data;
  - parton distributions constructed in a statistical physical picture of the nucleon:

$$p(x) \propto [\exp{(x - X_{0p})/\bar{x}} \pm 1]^{-1}$$

  - calculation done for A_1 at Q^2=5, 10, 100 (GeV/c)^2, ($g_2 = g_2^{ww}$)

World data on A_1^n and Soffer's statistical model



- Duality: W. Melnitchouk, Phys. Rev. Lett. **86**, 35 (2001)
  - local quark-hadron duality;
  - predict $x \sim 1$ behavior of A_1 from available data on G_E,G_M;

World data on A_1^n, Prediction from Duality



- Chiral Quark-Soliton Model:

  H.Weigel, L.Gamberg and H.Reinhardt, Phys. Lett. **B399**, 287 (1997), Phys. Rev. **D55**, 6910 (1997).

  – Two flavor chiral soliton approach to baryons, based on the Bosonized Nambu-Jona-Lasinio (NJL) model;

  – Model degrees of freedom are effective constituent quarks;

  – Tentative calculation done for $\frac{g_1}{F_1}$ at Q^2=1.5, 2.0, 3.0 (GeV/c)^2.

World data on A_1^n and chiral quark-soliton model



A_1^n World Data and Theoretical Predictions



Experiment E99-117

MEASURED A_1^n AT

x_{Bj}	0.331	0.474	0.609
Q^2 (GeV/c)^2	2.738	3.567	4.887
W^2 (GeV/c)^2	6.426	4.846	4.023

through e^- asymmetries $A_{\|}$ and $A_{\perp}$ in inclusive $\vec{e}^{\leftarrow} - {}^3\vec{H}e$ DIS;

EXPERIMENTAL SETUP

- Jefferson Lab(JLab) polarized e^- beam, 5.734 GeV, $P_{beam} = 81\%$, 12 μA
- Hall A polarized ${}^3He$ target, $\sim$ 10 atm, $P_{targ} = 40\%$
- Two Hall A High Resolution Spectrometers (HRS) at symmetric positions

THE JEFFERSON LAB ACCELERATOR



EXPERIMENTAL HALL A



- *Two VDCs* for tracking;
- *Sets of scintillators* for triggering on charged particles;
- *Gas cherenkov detector* for pion rejection;
- *Two layers of lead glass counters* for additional PID;
- *Pion rejection efficiency:* better than 10^{-4} with electron efficiency 99%.



Polarzied ^3He *Target*

PRINCIPLE OF POLARIZING

- Optical pumping of Rb by circularly polarized light



- Spin exchange through Rb-^3He collisions



THE CELL

- Cell density: 7.5∼10 amg.
- Maximum polarization: ∼40% (in beam) - NMR, EPR Polarimetries
- Cell lifetime: $\Gamma = \Gamma_{He} + \Gamma_{wall} + \Gamma_{beam} + \Gamma_{\Delta B}$
  $\sim (40)^{-1} = (744/[He])^{-1} \quad (90)^{-1} \quad (655/I)^{-1} \quad (1700)^{-1}$ (hours)$^{-1}$
- Cell end windows 100μm, wall thickness $\sim$ 130 mm
- Polarization diffusion: $P_p - P_t \sim 1\%$



TARGET SETUP



NMR POLARIMETRY (K.Kramer, PST2001)



EPR POLARIMETRY

- Rb Zeeman splitting $\propto$ Magnetic field magnitude $\vec{B}$;
- $P_{^3He}$ induce a small $\vec{B}_{^3He}$, $\vec{B} = \vec{B}_{^3He} + \vec{B}_{main}$;
- Measure Rb resonance frequency shift due to $P_{^3He}$



TARGET IN SITU



Polarized ^{3}He Target Performance During E99–117

Cell Name	Field direction	0°	180°	270°
Gore	June 06 ~ July 03	37%	35%	43%
Tilghman	July 13 ~ July 31	45%	43%	39%



Asymmetry Analysis

Electron asymmetries

$$A_{raw} = \frac{N^+/Q^+ - N^-/Q^-}{N^+/Q^+ + N^-/Q^-}$$

$$A_{\parallel,\perp} = \frac{A_{raw}}{f P_b P_t}$$

^{3}He asymmetries

$$A_1 = \frac{A_\parallel}{D(1+\eta\xi)} - \frac{\eta A_\perp}{d(1+\eta\xi)}$$

$$A_2 = \gamma\left\{\frac{1-\frac{y}{2}}{D'}A_\parallel + \frac{1}{D'}\left[\tan(\theta/2) + \frac{y}{2\sin\theta}\frac{1+(1-y)\cos\theta}{1-y}\right]A_\perp\right\}$$

$$g_1(x,Q^2) = \frac{F_1(x,Q^2)}{D'}\left[A_\parallel + \tan(\theta/2)\cdot A_\perp\right]$$

$$g_2(x,Q^2) = \frac{F_1(x,Q^2)}{2D'y\sin\theta}\left[\frac{E+E'\cos\theta}{E'}A_\perp - \sin\theta\cdot A_\parallel\right]$$

Asymmetries

^{3}He Raw Asymmetries



^{3}He Physics Asymmetries



From ^{3}He to Neutron

$^3\vec{He}$ AS EFFECTIVE $\vec{n}$



$\vec{^3He} \approx \vec{n}$

CONVOLUTION APPROACH

Ciofi degli Atti et.al., Phys.Rev.**C48**, 968(1993); Phys. Lett., **B404**, 223(1997)

- ^3He consists S, S', D
- Three body calculation using Fadeev wavefunction

$$\tilde{g_1^n} = \frac{1}{\rho_n}(g_1^{^3\mathrm{He}} - 2\rho_p g_1^p)$$

$$\tilde{A_1^n} = \frac{W_1^{^3\mathrm{He}}}{W_1^n}\frac{1}{\rho_n}(A_1^{^3\mathrm{He}} - 2\frac{W_1^p}{W_1^{^3\mathrm{He}}}\rho_p A_1^p)$$

COMPLETE ANALYSIS

F. Bissey et. al., hep-ph/0109069

- S, S', D, Δ isobar in ^3He wavefuntion

$$A_1^n = \frac{F_2^{^3He}}{P_n F_2^n(1+\frac{0.056}{P_n})}[A_1^{^3He} - 2\frac{F_2^p}{F_2^{^3He}}P_p A_1^p(1-\frac{0.014}{2P_p})]$$

Other Inputs

- $R(x, Q^2)$ - E143, K. Abe et. al., hep-ex/9808028
- F_2^p, F_2^D - NMC95, M. Arneodo et. al., hep-ph/9509406,
- EMC $F_2^{^3He} = \mathcal{R}^{^3He}(2F_2^p + F_2^n)$, $F_2^n = \frac{F_2^D}{\mathcal{R}^D} - F_2^p$, W. Melnitchouk, pri.comm.
- Effective nucleon polarization P_p, P_n
  - $P_p = -0.028 \pm 0.004$,$P_n = 0.86 \pm 0.02$
    C. Ciofi degli Atti et.al.,*Phys.Rev.***C48**, R968(1993)
  - $P_p = -0.021$,$P_n = 0.879$
    PEST5, F. Bissey et. al., *Phys. Rev.* **C64**, 024004(2001)
  - $P_p = -0.018$,$P_n = 0.897$
    CD-Bonn2000 + TM (Tucson-Melbourne) 3NF, A. Nogga, Ph.D. thesis, (2001)
- A_1^p - E155 fit, P. L. Anthony et. al., *Phys.Lett.***B493**:19-28,(2000) (hep-ph/0007248)
- g_2, using g_2^{ww} (E155x)

A_1^n Preliminary Result



A_1^n Preliminary Result



A_1^n *Preliminary Result*



Error Analysis

Statistics

- ΔA_1^n =5% at $x = 0.61$;

Experimental systematics

- Beam energy: 5.734 GeV, $\frac{\Delta E_b}{E_b} < 5 \times 10^{-4}$;
- Spectrometer momentum: $\frac{\Delta E_e}{E_e} < 5 \times 10^{-4}$;
- Spectrometer angle: $\Delta\theta_e < 0.1^\circ$;
- Beam polarization: 81%, $\frac{\Delta P_b}{P_b}$ <3%;
- Target polarization: 40%, $\frac{\Delta P_t}{P_t}$ <4%;
- Target spin orientation: $\Delta\theta_{targ}$ <1°;

Radiative Correction

- Correct $A_{\parallel}^{^3He}$, $A_{\perp}^{^3He}$ directly;
- F_2 variation (5 models), $F_2^{p,\mathrm{res}}$ (2);
- g_1 variation, g_1/F_1 variation (3), $g_1^{^3He,\mathrm{res}}$ variation;
- g_2 variation (g_2^{ww} vs. $g_2 = 0$);

Nuclear correction

- F_2^p, F_2^d; $\mathcal{R}^{^3He}$, $\mathcal{R}^D$; A_1^p; P_p, P_n

Error Analysis



Preliminary Result



black solid curve: g_1^n obtained from E155 fit for $\frac{g1}{F1}$ and NMC95 fit for F_1, at $Q^2 = 4.0$ (GeV/c)^2

Preliminary Result



Preliminary Result



red solid curve: xg_2^{ww} obtained from E155 fit for $\frac{g1}{F1}$ and NMC95 fit for F_1, at $Q^2 = 4.0$ (GeV/c)^2

Discussions

A_1^n and Spin Crisis

- Proton spin 'crisis': $\Sigma = \Sigma_v + \sum \Delta(q+\bar{q})_{sea} \sim 0.3 \pm 0.1$
- Current understanding of nucleon spin:
  - $\Sigma_v = 0.75$ from rCQM;
  - $\sum \Delta(q+\bar{q})_{sea} = -0.45$ from data;
  - $1 = \underbrace{\Sigma_v + \sum \Delta(q+\bar{q})_{sea}}_{\Sigma=0.3\pm0.1} + 2L_q + \Sigma_G$
- Spin physics programs have been planned at HERMES, RHIC and CERN to study $\Delta\bar{q}$ and ΔG;
- Measuring $\Delta\bar{q}$ and ΔG is also based on the knowing of contribtuion from valence quarks;
- A_1^n and A_1^p at large x will check our valence quark picture of $\Sigma_v = 0.75$. A negative A_1^n will bring a great doubt at this assumption and thus start a new search for resolution of 'spin crisis'.

- Can also check pQCD at valence quark region. ($\frac{\Delta d}{d} \rightarrow$? 1, $-\frac{1}{3}$)

Summary

EXPERIMENT E99-117

- Provides the first precise data of A_1^n and g_1^n at $x > 0.4$;
- Data on A_2^n, g_2^n also available;
- Check current understanding of nucleon spin;
- Help to understand constituent quark models;
- Provide new constraint on other models, PDFs.

Summary

NEUTRON SPIN STRUCTURE PROGRAM AT JLAB HALL A (USING POLARIZED ^3He)

Sum Rules

- ● E94010 A Connection between Bjorken and GDH Sum Rules, (Nov∼Dec 1998)
- ○ E97110 The Generalized GDH Sum Rule for Nearly Real Photons (Sep 2002)

Spin Structure Functions

- ● E99117 Neutron Asymmetry A_1^n at larget x_{Bj} (June $\sim$ July 2001)
- ● E97103 Search for Higher Twist Effect in g_2^n (Aug $\sim$ Sep 2001)
- ○ E01012 Neutron Spin Structure in the Resonance Region (approved)

A_1^n @ JLab 12 GeV Upgrade

- Precision measurement of spin asymmetry in valence quark region
- Decisive test of pQCD vs quark model, insight to quark-gluon wavefunctions