Precision Measurement of the Neutron Spin Dependent Structure Functions

A Dissertation Presented

by

Yury G. Kolomensky

Submitted to the Graduate School of the
University of Massachusetts Amherst in partial fulfillment
of the requirements for the degree of

Doctor of Philosophy

February 1997

Department of Physics and Astronomy

PRECISION MEASUREMENT OF THE NEUTRON
SPIN DEPENDENT STRUCTURE FUNCTIONS

A Dissertation Presented

by

YURY G. KOLOMENSKY

Approved as to style and content by:

Gerald A. Peterson, Chair

Ross S. Hicks, Member

Eugene Golowich, Member

Gerassimos G. Petratos, Member

Michael F. Skrutskie, Member

John F. Donoghue, Department Head
Department of Physics and Astronomy

Оле, Мише и Саше

Acknowledgments

First and foremost, I would like to express my gratitude to my adviser Professor Gerald A. Peterson for giving me the opportunity to work on this experiment from its very beginning in 1994. I am deeply indebted to him for his attention, help, and advice I value very much. I would like to thank the members of my committee for taking the time to read and correct the manuscript.

The experiment was a collaborative effort of many physicists. I would like to thank the spokesperson, Professor Emlyn W. Hughes, for the time and energy he put into the project, and for being such an enthusiastic and motivating person he is. I had a privilege to work with many fine physicists, and I appreciate help, support, and many discussions with Ray Arnold, Cristina Berisso, Gordon Cates, Roger Gearhart, Krishna Kumar, Zein-Eddine Meziani, Makis Petratos, Rainer Pitthan, Richard Prepost, Paul Souder, and Charlie Young. This work would not be possible without the effort of my fellow graduate students. Many thanks to Piotr, MO, Mike, Franck, Dave, Todd, Greg, Steve, Sebastien, and Jason for sharing this exciting experience with me, and for making the project succeed.

My physics education started at St. Petersburg Technical University in Russia, and I am thankful to the faculty of the Department of Physics and Mechanics for giving me the basis for a successful career in physics. I was fortunate to continue my education at UMass and interact with Professors Eugene Golowich, John Donoghue, and Barry Holstein.

Finally, I would like to thank Alexey Petrov for his friendship and help, and my parents for their support and understanding. But most of all, I am indebted to my dearest wife Olga for her love, patience, and support through all these years.

Abstract

Precision Measurement of the Neutron Spin Dependent Structure Functions

February 1997

Yury G. Kolomensky, B.S., St. Petersburg Technical University, Russia

M.S., University of Massachusetts Amherst

Ph.D., University of Massachusetts Amherst

Directed by: Professor Gerald A. Peterson

In experiment E154 at the Stanford Linear Accelerator Center the spin dependent structure function $g_1^n(x, Q^2)$ of the neutron was measured by scattering longitudinally polarized 48.3 GeV electrons off a longitudinally polarized ^{3}He target. The high beam energy allowed us to extend the kinematic coverage compared to the previous SLAC experiments to $0.014 \leq x \leq 0.7$ with an average Q^2 of 5 GeV^2. We report the integral of the spin dependent structure function in the measured range to be $\int_{0.014}^{0.7} dx\ g_1^n(x, 5\ \mathrm{GeV}^2) = -0.036 \pm 0.004(\mathrm{stat.}) \pm 0.005(\mathrm{syst.})$. We observe relatively large values of g_1^n at low x that call into question the reliability of data extrapolation to $x \to 0$. Such divergent behavior disagrees with predictions of the conventional Regge theory, but is qualitatively explained by perturbative QCD. We perform a Next-to-Leading Order perturbative QCD analysis of the world data on the nucleon spin dependent structure functions g_1^p and g_1^n paying careful attention to the experimental and theoretical uncertainties. Using the parameterizations of the helicity-dependent parton distributions obtained in the analysis, we evolve the data to $Q^2 = 5$ GeV^2, determine the first moments of the polarized structure functions of the proton and neutron, and find agreement with the Bjorken sum rule.

Table of Contents

List of Tables

List of Figures

CHAPTER 1

MOTIVATION

1.1 Introduction

The mystery of the structure of the nucleon is one of the most fascinating challenges facing modern physics. Deep inelastic scattering data helps us to meet the challenge. Like a powerful microscope, it allows us to look inside the sub-nuclear particles at a much deeper level and check our ideas of what nucleons consist of and how they behave.

Deep-inelastic scattering (DIS) experiments provide perhaps our cleanest window on hadron structure at large momentum transfer squared Q^2. The original DIS experiments in End Station A at the Stanford Linear Accelerator Center (SLAC) in the 1960s showed that the form factors in ep scattering exhibit approximate *scaling* at large Q^2.[1] This remarkable observation was celebrated when the 1990 Nobel Prize in Physics was awarded to the experimenters Friedman, Kendall, and Taylor.[2] It gave rise to the original parton ideas of Feynman[3] and of Bjorken and Paschos.[4,5] Precise data revealed the logarithmic scaling violations and also the gluon distribution that were predicted by QCD.

While the theoretical grounds for understanding unpolarized deep inelastic scattering are well established and surprises are rare, experimental polarization data have often challenged the theory. It is more than two decades since the first polarized deep inelastic scattering experiments were done at SLAC.[6,7] This work became the subject of renewed interest when EMC[8] extended the SLAC measurements to smaller x and announced their results on the spin-dependent proton structure function g_1. The naive parton model interpretation of their data is that the quarks contribute very little to the proton's spin – in contradiction with quark models.

Since that measurement, much more data have become available from SLAC[9–11] and CERN.[12,13] The work described in these pages is a further step in studying the dynamics of quarks and gluons inside the nucleon. In experiment E154,[14] we have measured the spin structure functions of the neutron $g_1^n(x, Q^2)$ and $g_2^n(x, Q^2)$ by scattering longitudinally polarized electrons off longitudinally or transversely polarized ^{3}He nuclei. High statistical precision and broad kinematic coverage of the experiment gives us a stronger basis for understanding the structure of the nucleon.

The experiment was carried out in October-November of 1995 by a collaboration of 80 physicists from 23 institutions in End Station A at SLAC.

1.2 Structure functions

1.2.1 Definitions

Our primary concern is with deep-inelastic scattering from polarized targets. Experiments at SLAC and DESY use polarized electrons while experiments at CERN by the European Muon Collaboration (EMC) and the Spin Muon Collaboration (SMC) utilize naturally polarized muon beams. Consider a DIS experiment where a lepton beam with definite polarization and momentum $k^\mu = (E, \vec{k})$ scatters from a polarized proton target. This is shown in the one photon exchange diagram of Fig. 1.1 to leading order in the electromagnetic interaction. We work in the laboratory frame so that the proton target P has momentum $p^\mu = (M, 0)$ and polarization S^μ. The lepton L is scattered through an angle θ and emerges with momentum $k'^\mu = (E', \vec{k}')$. The exchanged photon carries four-momentum $q^\mu = (k - k')^\mu$. The scattering process is then characterized by the two invariants $Q^2 = -q^2$ and $\nu = p \cdot q/M$ [$\nu = (E - E')$ in the LAB frame] or, equivalently, by Q^2 and the Bjorken variable $x = \frac{Q^2}{2M\nu}$. We measure the inclusive hadronic cross section so that hadronic final states X with



Fig. 1.1. Deep inelastic scattering.

the same invariant mass squared, $W^2 = (p+q)^2$, are not separated. The kinematic variables are summarized in Table 1.1.

The differential cross section for the one photon exchange process (Fig. 1.1) is given by[15]

$$\frac{d^2\sigma}{d\Omega dE'} = \frac{\alpha^2}{Q^4}\frac{E'}{E} L_{\mu\nu} W^{\mu\nu}. \tag{1.1}$$

Here, α is the fine structure constant and $L_{\mu\nu}$ and $W_{\mu\nu}$ describe the leptonic and hadronic vertices respectively. Since the lepton is elementary we can write down an exact expression for $L_{\mu\nu}$ from the Feynman rules,

$$L_{\mu\nu} \equiv L^S_{\mu\nu} + iL^A_{\mu\nu} = 2\left[k_\mu k'_\nu + k'_\mu k_\nu - g_{\mu\nu}(k \cdot k' - m^2) + im\epsilon_{\mu\nu\rho\sigma} q^\rho s^\sigma\right] \tag{1.2}$$

where we split the leptonic tensor in two parts: $L^S_{\mu\nu}$ is symmetric under $\mu \Leftrightarrow \nu$, and $L^A_{\mu\nu}$ is antisymmetric and contains the lepton spin.

The hadronic tensor $W^{\mu\nu}$ contains all of the information about the hadronic target that one can extract from such inclusive measurements. Its form is constrained by symmetry arguments. Again, we write $W_{\mu\nu}$ as a sum of symmetric and antisym-

Table 1.1. Kinematic variables and invariants of lepton-nucleon scattering.

Variable	Expression	Lab Frame	Meaning
E			Incident lepton energy
E'			Scattered lepton energy
$\vec{k}$			Incident lepton momentum
$\vec{k}'$			Scattered lepton momentum
ϑ	$\arccos\left(\frac{\vec{k}\cdot\vec{k}'}{\lvert\vec{k}\rvert\lvert\vec{k}'\rvert}\right)$		Lepton scattering angle
m			Lepton mass
k^μ	$(E,\vec{k})$		Incident lepton 4-momentum
k'^μ	$(E',\vec{k}')$		Scattered lepton 4-momentum
s^μ			Incident lepton polarization vector
M			Nucleon mass
P^μ		$(M,\vec{0})$	Nucleon 4-momentum
S^μ			Nucleon polarization vector
q^μ	$(k-k')^\mu$		4-momentum transfer
Q^2	$-q^2$	$4EE'sin^2\vartheta/2$	4-momentum transfer squared
ν	$P\cdot q/2M$	$(E-E')$	Energy transfer
x	$Q^2/2M\nu$	$\frac{4EE'sin^2(\vartheta/2)}{2M(E-E')}$	Bjorken x;
			the fraction of nucleon momentum carried by a struck parton

metric contributions; $W_{\mu\nu} = W^S_{\mu\nu} + iW^A_{\mu\nu}$. Then the requirements of covariance, parity, charge conjugation, and current conservation ($q^\mu W_{\mu\nu} = 0$) imply the form

$$W^S_{\mu\nu} = \frac{1}{M}(-g_{\mu\nu} + \frac{q_\mu q_\nu}{q^2})F_1(x,Q^2) + \frac{1}{M^2\nu}(p_\mu - \frac{p\cdot q}{q^2}q_\mu)(p_\nu - \frac{p\cdot q}{q^2}q_\nu)F_2(x,Q^2) \quad (1.3)$$

and

$$W^A_{\mu\nu} = \frac{\epsilon_{\mu\nu\lambda\sigma}q^\lambda S^\sigma}{M\nu}g_1(x,Q^2) + \frac{\epsilon_{\mu\nu\lambda\sigma}q^\lambda(M\nu S^\sigma - q\cdot Sp^\sigma)}{M^2\nu^2}g_2(x,Q^2) \quad (1.4)$$

The form factors in Equations 1.3 and 1.4 contain all of the target dependent information.

One has to use both polarized beam and polarized target in order to measure spin-dependent structure functions g_1 and g_2 since combinations with mixed symmetry vanish $L^S_{\mu\nu}W^{\mu\nu}_A = L^A_{\mu\nu}W^{\mu\nu}_S = 0$ and only combinations $L^S_{\mu\nu}W^{\mu\nu}_S$ and $L^A_{\mu\nu}W^{\mu\nu}_A$

survive. For this experiment, the unpolarized structure functions F_1 and F_2 are considered to be known.

Deep inelastic scattering is defined by the kinematic limit where both $Q^2 \gg M^2$ and $W^2 \gg M^2$, so that we are beyond the resonance region (where W^2 may coincide with the mass squared of one of the excited nucleon resonances). In the DIS limit the form factors F_i and g_i in $W_{\mu\nu}$ exhibit approximate *scaling*. That is, they behave as structure functions of the single variable x - modulo a slow logarithmic variation in Q^2, which is described by perturbative QCD. The scaling property reveals a local interaction between the hard photon and charged elementary partons (quarks) inside the proton. It is the same effect as in Rutherford's α particle scattering experiments which revealed the nucleus inside the atom.[16]

We now turn to the measurement of the spin dependent structure functions. We will consider leptons longitudinally polarized and targets longitudinally or transversely polarized with respect to the incident beam direction. We will let $\uparrow\downarrow$ denote the longitudinal lepton polarization and $\Uparrow\Downarrow$ ($\Leftarrow$) denote longitudinal (transverse) polarization of the target nucleon. Then the differential cross sections are

$$\frac{d^2\sigma}{d\Omega dE'}^{\downarrow\Uparrow} + \frac{d^2\sigma}{d\Omega dE'}^{\uparrow\Uparrow} = \frac{8\alpha^2(E')^2}{MQ^4}\Big[2\sin^2\frac{\vartheta}{2}\ F_1(x,Q^2) + \frac{M}{\nu}\cos^2\frac{\vartheta}{2}\ F_2(x,Q^2)\Big] \quad (1.5)$$

and

$$\frac{d^2\sigma}{d\Omega dE'}^{\downarrow\Uparrow} - \frac{d^2\sigma}{d\Omega dE'}^{\uparrow\Uparrow} = \frac{4\alpha^2 E'}{Q^2 EM\nu}\Big[(E + E'\cos\vartheta)\ g_1(x,Q^2) - 2xM\ g_2(x,Q^2)\Big] \quad (1.6)$$

for longitudinally polarized leptons and nucleons, and

$$\frac{d^2\sigma}{d\Omega dE'}^{\downarrow\Leftarrow} - \frac{d^2\sigma}{d\Omega dE'}^{\uparrow\Leftarrow} = \frac{4\alpha^2 E'}{Q^2 EM\nu}E'\sin\vartheta\Big[\ g_1(x,Q^2) + \frac{4xEM}{Q^2}\ g_2(x,Q^2)\Big] \quad (1.7)$$

for longitudinally polarized leptons and transversely polarized targets.

The structure function g_2 is suppressed in Eq. (1.6) with respect to g_1 by a factor $\frac{M}{E} \approx 0.02$, for the beam energy of 50 GeV. The transverse asymmetry in Eq. (1.7) primarily measures g_2.

The experiments usually do not measure cross section differences of Equations (1.6) and (1.7) directly, but measure longitudinal and transverse asymmetries

$$A_{\|} = \frac{\frac{d^2\sigma}{d\Omega dE'}^{\downarrow\Uparrow} - \frac{d^2\sigma}{d\Omega dE'}^{\uparrow\Uparrow}}{\frac{d^2\sigma}{d\Omega dE'}^{\downarrow\Uparrow} + \frac{d^2\sigma}{d\Omega dE'}^{\uparrow\Uparrow}} \tag{1.8}$$

and

$$A_{\perp} = \frac{\frac{d^2\sigma}{d\Omega dE'}^{\downarrow\Leftarrow} - \frac{d^2\sigma}{d\Omega dE'}^{\uparrow\Leftarrow}}{\frac{d^2\sigma}{d\Omega dE'}^{\downarrow\Leftarrow} + \frac{d^2\sigma}{d\Omega dE'}^{\uparrow\Leftarrow}}. \tag{1.9}$$

Then, if the beam and/or target polarizations are reversed frequently, all slow-varying factors (beam properties, detector efficiencies and acceptance, etc.) cancel between different polarization states, and one can write the asymmetries in terms of detector counting rates:

$$A_{\|} = \frac{1}{fP_bP_t}\frac{(N/Q)^{\downarrow\Uparrow} - (N/Q)^{\uparrow\Uparrow}}{(N/Q)^{\downarrow\Uparrow} + (N/Q)^{\uparrow\Uparrow}} \tag{1.10}$$

and

$$A_{\perp} = \frac{1}{fP_bP_t}\frac{(N/Q)^{\downarrow\Leftarrow} - (N/Q)^{\uparrow\Leftarrow}}{(N/Q)^{\downarrow\Leftarrow} + (N/Q)^{\uparrow\Leftarrow}}\,. \tag{1.11}$$

Here N and Q are the number of events and beam charge for each helicity state, respectively, P_b and P_t are the beam and target polarizations, and the factor f is the dilution factor, the ratio of scattering rates from the polarized nuclei to the total number of target nuclei:

$$f(x,Q^2) = \frac{n_P\sigma_P(x,Q^2)}{n_P\sigma_P(x,Q^2) + n_U\sigma_U(x,Q^2)}. \tag{1.12}$$

The polarized nucleus (^{3}He, H, or D) is denoted by P, and U stands for unpolarized material. The scattering cross sections are $\sigma_i(x,Q^2)$, and n_i are the nucleon densities.

1.2.2 The virtual photon-nucleon asymmetries

Deep inelastic scattering is composed of two processes. One, represented by the top vertex in Fig. 1.1, is the emission of the virtual photon by the incident lepton, and is completely calculable in QED. The bottom vertex represents the absorption of the photon by the target nucleon. The optical theorem relates the total photoabsorption cross section to the imaginary ("absorptive") part of the forward ($t = 0$) virtual photon-nucleon Compton scattering amplitude (where t is the momentum transfer Mandelstam variable).

Let us denote the Compton amplitudes $\mathcal{M}_{ab\to cd}$ where a and b are the incident photon and nucleon, and c and d are the final photon and nucleon helicities, respectively. These amplitudes are derived from the hadronic tensor $W_{\mu\nu}$ through

$$\mathcal{M}_{ab\to cd} = \epsilon^{\mu\dagger}(\lambda_a)\epsilon^{\nu}(\lambda_c)W_{\mu\nu} \ , \tag{1.13}$$

where $\epsilon^{\mu}(\lambda)$ is the polarization vector of the virtual photon.

There are four independent helicity amplitudes (just as there are four independent structure functions defined in Section 1.2.1):[17]

$$\mathcal{M}_{1\frac{1}{2}\to 1\frac{1}{2}},\ \mathcal{M}_{1-\frac{1}{2}\to 1-\frac{1}{2}},\ \mathcal{M}_{0\frac{1}{2}\to 0\frac{1}{2}},\ \mathcal{M}_{0\frac{1}{2}\to 1-\frac{1}{2}}. \tag{1.14}$$

Using Equations (1.3) and (1.4), we get for the virtual photoabsorption cross sections:

$$\begin{aligned}
\sigma_{1/2} &= \frac{4\pi^2\alpha}{K}\mathcal{M}_{1-\frac{1}{2}\to 1-\frac{1}{2}} = \frac{4\pi^2\alpha}{MK}\left[F_1 + g_1 - \frac{2Mx}{\nu}g_2\right] \\
\sigma_{3/2} &= \frac{4\pi^2\alpha}{K}\mathcal{M}_{1\frac{1}{2}\to 1\frac{1}{2}} = \frac{4\pi^2\alpha}{MK}\left[F_1 - g_1 + \frac{2Mx}{\nu}g_2\right] \\
\sigma_{L} &= \frac{4\pi^2\alpha}{K}\mathcal{M}_{0\frac{1}{2}\to 0\frac{1}{2}} = \frac{4\pi^2\alpha}{K}\left[\frac{F_2}{\nu}\left(1 + \frac{\nu^2}{Q^2}\right) - \frac{1}{M}F_1\right] \\
\sigma_{TL} &= \frac{4\pi^2\alpha}{K}\mathcal{M}_{0\frac{1}{2}\to 1-\frac{1}{2}} = \frac{4\pi^2\alpha}{K}\frac{\sqrt{Q^2}}{M\nu}\left[g_1 + g_2\right]
\end{aligned} \tag{1.15}$$

where the photon flux K is defined according to Hand's convention as

$$K = \frac{W^2 - M^2}{2M} = \nu - \frac{Q^2}{2M}. \tag{1.16}$$

The quantities $\sigma_{1/2}$ and $\sigma_{3/2}$ are the virtual photoabsorption cross sections when the total photon-nucleon spin along the photon direction is 1/2 and 3/2 respectively. The total transverse photoabsorption cross section is defined by

$$\sigma_T = \frac{1}{2}\left(\sigma_{1/2} + \sigma_{3/2}\right) = \frac{4\pi^2\alpha}{MK}F_1. \tag{1.17}$$

The cross section for the longitudinally polarized photon is σ_L, and the interference term between transverse and longitudinal amplitudes is σ_{TL}.

The virtual photon-nucleon asymmetries are defined by

$$A_1(x,Q^2) \equiv \frac{\sigma_{1/2}-\sigma_{3/2}}{\sigma_{1/2}+\sigma_{3/2}} = \frac{g_1(x,Q^2)-\gamma^2 g_2(x,Q^2)}{F_1(x,Q^2)} \tag{1.18}$$

and

$$A_2(x,Q^2) \equiv \frac{\sigma_{TL}}{\sigma_T} = \frac{\gamma\left(g_1(x,Q^2)+g_2(x,Q^2)\right)}{F_1(x,Q^2)}\ , \tag{1.19}$$

where $\gamma = \sqrt{Q^2/\nu^2}$. We can also define the ratio of the longitudinal to transverse cross sections

$$R(x,Q^2) \equiv \frac{\sigma_L}{\sigma_T} = \frac{1+\gamma^2}{2xF_1(x,Q^2)}F_2(x,Q^2) - 1 \tag{1.20}$$

which links two unpolarized structure functions

$$F_1(x,Q^2) = \frac{1+\gamma^2}{2x\left(1+R(x,Q^2)\right)}F_2(x,Q^2). \tag{1.21}$$

The virtual asymmetries satisfy the unitarity conditions[18]

$$|A_1(x,Q^2)| \leq 1, \quad |A_2(x,Q^2)| \leq \sqrt{R(x,Q^2)}. \tag{1.22}$$

1.2.3 Structure functions and experimental asymmetries

Finally, we can express the structure functions and virtual photon-nucleon asymmetries in terms of the experimentally measured asymmetries

$$g_1(x,Q^2) = \frac{F_1(x,Q^2)}{D'}\left[A_{||}(x,Q^2)+A_{\perp}(x,Q^2)\tan(\vartheta/2)\right]\ , \tag{1.23}$$

$$g_2(x,Q^2) = \frac{F_1(x,Q^2)}{D'}\frac{y}{2\sin\vartheta}\left[-A_{||}(x,Q^2)\sin\vartheta + A_{\perp}(x,Q^2)\frac{E+E'\cos\vartheta}{E'}\right] \tag{1.24}$$

and

$$A_1(x,Q^2) = \frac{A_{||}(x,Q^2)}{D(1+\zeta\eta)} - \frac{\eta A_{\perp}(x,Q^2)}{d(1+\zeta\eta)}\ , \tag{1.25}$$

$$A_2(x,Q^2) \;=\; \frac{\zeta A_{\|}(x,Q^2)}{D(1+\zeta\nu)} + \frac{A_{\perp}(x,Q^2)}{d(1+\zeta\nu)} \,. \qquad (1.26)$$

The kinematic factors used above are

$$\begin{aligned} \epsilon &= \frac{1}{1+2(1+(\nu^2/Q^2))\tan^2(\vartheta/2)}\,, \\ D' &= \frac{(1-\epsilon)(2-y)}{y(1+\epsilon R(x,Q^2))}\,, \\ D &= \frac{1-E'\epsilon/E}{1+\epsilon R(x,Q^2)}\,, \\ d &= D\sqrt{\frac{2\epsilon}{1+\epsilon}}\,, \\ \eta &= \frac{\epsilon\sqrt{Q^2}}{E-E'\epsilon}\,, \quad \text{and} \\ \zeta &= \eta\frac{1+\epsilon}{2\epsilon}\,, \end{aligned} \qquad (1.27)$$

where $y = (E-E')/E$ is the fractional energy transfer from the electron to hadrons.

1.2.4 The parton model and polarized parton distributions

The parton model began with Feynman who showed that the early SLAC deep inelastic scattering experiments could be explained in terms of the hard photon scattering incoherently from elementary *parton* constituents in the proton. The structure functions measure the probability for finding a quark with momentum fraction $x = p^+_{parton}/p^+_{proton}$ [1] in the proton and which is polarized either in the same or the opposite direction to the proton's polarization. This is usually called the naive parton model; it has no gauge degrees of freedom.

In the naive parton model the structure functions are described by the four linearly independent parton distributions. There are two spin-independent distribu-

[1] Here, p^+ denotes the light-cone momentum; see Appendix A for details

tions (one for a quark and one for an anti-quark), and two spin-dependent distributions. The charge conjugation even combination [2]

$$(q+\bar{q})^{\uparrow}(x)+(q+\bar{q})^{\downarrow}(x) \tag{1.28}$$

occurs in the structure functions $F_1(x,Q^2)$ and $F_2(x,Q^2)$. It is measured in the unpolarized lepton-nucleon scattering. In the naive parton model one finds

$$F_2(x)=2xF_1(x)=x\sum_q e_q^2\Big[(q+\bar{q})^{\uparrow}(x)+(q+\bar{q})^{\downarrow}(x)\Big], \tag{1.29}$$

where e_q denotes the quark charge. The relationship between F_1 and F_2 is called the Callan-Gross relation.

The polarized (spin-dependent) quark and anti-quark distributions occur in the spin-dependent structure function $g_1(x)$, which in the naive parton model is written as:

$$g_1(x)=\frac{1}{2}\sum_q e_q^2\Delta q(x)\ , \tag{1.30}$$

where

$$\Delta q(x)\equiv\delta q(x)+\delta\bar{q}(x)=(q^{\uparrow}-q^{\downarrow})(x)+(\bar{q}^{\uparrow}-\bar{q}^{\downarrow})(x) \tag{1.31}$$

is the polarized quark distribution. It is helpful to rewrite $g_1(x)$ in terms of the SU(3) flavor combinations:

$$\begin{aligned}\Delta q_3(x) &\equiv \Delta u(x)-\Delta d(x),\\ \Delta q_8(x) &\equiv \Delta u(x)+\Delta d(x)-2\Delta s(x),\\ \Delta\Sigma(x) &\equiv \Delta u(x)+\Delta d(x)+\Delta s(x).\end{aligned} \tag{1.32}$$

Combination $\Delta\Sigma$ (also refered to as Δq_0) is a singlet distribution whereas Δq_3 and Δq_8 are non-singlet.

[2] In Eq. (1.28), $q^{\uparrow}(x)$ ($\overline{q}^{\uparrow}(x)$) denotes the distribution for helicity aligned quark (anti-quark), and $q^{\downarrow}(x)$ ($\overline{q}^{\downarrow}(x)$) stands for helicity anti-aligned distributions. By convention, the parton distributions are given for the proton, and one can get the corresponding neutron distribution functions using isospin symmetry. These distributions are often written in the literature as $G_{p/H}(x,\lambda,Q)$ where p denotes the parton (quark or gluon), H stands for the hadron, and λ is the parton helicity.

Calculation of the x-dependence of the structure functions requires precise knowledge of the nucleon wave function. The moments of structure functions, on the other hand, can be calculated and provide a testing ground for our understanding of nucleon structure in QCD. The first piece of experimental evidence that showed that we needed to go beyond the naive parton model came when it was observed that quarks contribute only 50% of the proton's momentum. This was naturally accommodated with the advent of QCD where the gluons in the limit of infinite Q^2 carry the momentum fraction

$$\langle xG_{g/p}(x,Q^2)\rangle \to \frac{16}{16+3N_f} \, , \tag{1.33}$$

where N_f is the number of quark flavors.[19]

In the naive parton model,

$$\Delta q = \int_0^1 dx \ \Delta q(x) \tag{1.34}$$

determines the fraction of the proton's helicity which is carried by quarks (and anti-quarks) of flavor q. Thus, we write

$$\Gamma_1^p \equiv \int_0^1 dx \ g_1^p(x) = \frac{1}{12}(\Delta u - \Delta d) + \frac{1}{36}(\Delta u + \Delta d - 2\Delta s) + \frac{1}{9}(\Delta u + \Delta d + \Delta s) + \frac{2}{9}\Delta c. \tag{1.35}$$

Assuming isospin symmetry, the neutron integral is obtained by interchanging u and d quarks in Eq. (1.35). The singlet term $\Delta\Sigma = \Delta u + \Delta d + \Delta s$ denotes the light-quark spin, or more strictly speaking, helicity content of the nucleon[3]

In operator language, Δq is defined by the proton matrix element of the axial current. We write

$$2MS_\mu \Delta q_a = < p, S|\bar{q}\gamma_\mu\gamma_5 \frac{\lambda^a}{2} q|p, S > \tag{1.36}$$

for $a = 3, 8, 0$. The non-singlet matrix elements also arise in the neutron and hyperon beta decays. Current algebra relates the spin dependent (strong interaction) struc-

[3] In Eq. (1.35), we have explicitly written a contribution from charm quarks Δc, which is present above the charm threshold. It is usually ignored in the numerical analysis.

ture of the proton measured in polarized deep inelastic scattering at high energies to the quantities needed in low-energy weak-interaction physics. The currents which measure Δq_3 and Δq_8 do not renormalize, so these quantities are scale independent. They are determined as $\Delta q_3 = g_A = F + D$ and $\Delta q_8 = \frac{1}{\sqrt{3}}(3F - D)$ within SU(3). The axial coupling constant of the neutron beta-decay is g_A, and F and D are the antisymmetric and symmetric SU(3) couplings. One finds[20,21]

$$\Delta u - \Delta d = g_A = 1.2601 \pm 0.0025 \tag{1.37}$$

and

$$\Delta u + \Delta d - 2\Delta s = \frac{1}{\sqrt{3}}(3F - D) = 0.688 \pm 0.035. \tag{1.38}$$

Since Δq_3 and Δq_8 are determined from other experiments, by measuring the integral Γ_1^p or Γ_1^n, we can extract the singlet "spin content" of the proton $\Delta\Sigma$ as well as individual quark contributions Δu, Δd, and Δs (this assumes that there is a negligible charm component Δc in the data.)

1.2.5 Sum rules

The angular momentum of a fast-moving nucleon has three sources, the angular momentum carried by the quarks, the angular momentum carried by the gluons, and the orbital angular momentum carried by any of the constituents. Angular momentum conservation for J_z at a fixed light-cone time implies the sum rule[24]4

$$\frac{1}{2}(\Delta u + \Delta d + \Delta s) + \Delta G + < L_z >= \frac{1}{2}. \tag{1.39}$$

The sum $\Delta\Sigma = \Delta u + \Delta d + \Delta s$ is interpreted as the proton helicity carried by quarks, and $\Delta G \equiv \int_0^1 dx \Delta G(x)$ is the helicity carried by the gluon where $\Delta G(x)$ is the difference between the helicity-aligned and anti-aligned gluon distributions

4Such interpretation is not gauge- or Lorenz-invariant[25], so one commonly assumes the infinite-momentum frame (in which the nucleon moves with an infinite momentum along the z direction), and the light-cone gauge (see Appendix A). A gauge-invariant treatment of the spin sum rule has been derived in Ref. [25].

$G^+(x)$ and $G^-(x)$. The unpolarized gluon distribution $G(x)$ is the sum of these two functions, $G(x) \equiv G^+(x) + G^-(x)$. By definition, the anti-quark contributions are included in $\Delta q(x)$ and $q(x)$.

There are two sum rules for g_1 which can be tested in spin dependent deep inelastic scattering. The Bjorken sum rule[26] gives a relation for the difference between the first moment of g_1 for a proton and neutron target. In the scaling limit, it reads:

$$\int_0^1 dx\ \Big(g_1^p(x) - g_1^n(x)\Big) = \frac{1}{6} g_A\ . \tag{1.40}$$

The Bjorken sum rule was derived using current algebra before the advent of QCD and is a test of isospin symmetry.

At the finite Q^2 of an experiment one must include perturbative QCD (pQCD) Wilson coefficients. Including corrections to order α_S^3, the Bjorken sum rule becomes

$$\int_0^1 dx\ \Big(g_1^p(x) - g_1^n(x)\Big) = \frac{1}{6} g_A \Big[1 - \frac{\alpha_S(Q^2)}{\pi} + C_2 \left(\frac{\alpha_S(Q^2)}{\pi}\right)^2 + C_3 \left(\frac{\alpha_S(Q^2)}{\pi}\right)^3\Big], \tag{1.41}$$

where $C_2 = -3.5833$ and $C_3 = -20.2153$ in the three-flavor theory.[28] The corresponding expansion for the first moment of the proton spin structure function reads

$$\int_0^1 dx\ g_1^p(x, Q^2) = \frac{1}{6}\Big[\left(\Delta q_3 + \frac{1}{\sqrt{3}}\Delta q_8\right)\left(1 - \frac{\alpha_S}{\pi}\right) + 2\sqrt{\frac{2}{3}}\Delta\Sigma\left(1 - \frac{\alpha_S}{\pi}\frac{33 - 8N_f}{33 - 2N_f}\right)\Big], \tag{1.42}$$

where the perturbative coefficients are quoted to $O(\alpha_S)$.[27]

The second sum rule for g_1 is the Ellis-Jaffe sum rule[29] and is a test of Zweig's (or OZI after Okuba, Zweig, and Iizuka) rule in the flavor singlet channel. If we assume that strange (and heavy) quarks do not play a significant role and set $\Delta s = 0$, then the quark "spin content" would be determined by the hyperon beta decays. In this scenario we would have $\Delta\Sigma = \Delta q_8 = 0.688 \pm 0.035$, where the rest of the proton's spin would be carried by the gluons and also by quark and gluon orbital angular momentum. Substituting this value into Eq. (1.42) we find

$$\int_0^1 dx\ g_1^p(x) = 0.189 \pm 0.005 \tag{1.43}$$

(at 10 GeV^2). The Ellis-Jaffe sum rule involves a model-dependent assumption that the OZI rule is obeyed, whereas the Bjorken sum rule should hold exactly in QCD. Experiments up to now have found Ellis-Jaffe sum rule to be violated by more than two standard deviations[8,10–13] and confirmed the Bjorken sum rule.[11,13,30]

1.2.6 Transverse spin-dependent structure function g_2

The nucleon transverse spin-dependent structure function $g_2(x, Q^2)$ has only recently been measured,[22,23] and there have been few theoretical studies of it. As we mentioned above, it is strongly suppressed in the longitudinal asymmetry measurements and has been customarily neglected in the analysis of the early experiments. However, precise measurements of g_1, like the one described in this thesis, require taking into account all associated corrections.

One of the first theoretical observations was the Burkhardt-Cottingham sum rule[31]

$$\int_0^1 dx\ g_2(x, Q^2) = 0\ . \tag{1.44}$$

It was originally derived using Regge theory and dispersion relations,[32] raised many questions and induced a lively theoretical discussion.[32–34] The experimental measurement of the sum rule could provide a test of Regge theory in DIS.

A reliable method for exploring the properties of structure functions in the DIS limit is the operator product expansion (OPE) on the light-cone, *i.e.* in the space-time region $x_\mu x^\mu \leq Q^{-2} \to 0$ in the Bjorken limit.[32] Using OPE, the product of two electromagnetic currents forming the hadronic tensor

$$W_{\mu\nu}(q,p,S) = \frac{1}{4\pi}\int d^4x e^{iqx}\langle pS|[J_\mu(x), J_\nu(0)]|pS\rangle \tag{1.45}$$

can be expanded in terms of the renormalized local operators with singular coefficients. In the Bjorken limit the importance of an operator is determined by its twist, or light-cone singularity of the coefficient function (see Section 1.3.2.2). As it is shown in Ref. [35], while higher-twist operators (for instance, twist-three operators that reflect finite quark masses and quark-gluon interactions in the nucleon) have only correcting impact on g_1, their contribution to g_2 is rather big and can be measured experimentally. In general, g_2 can be written as

$$g_2(x,Q^2) = g_2^{WW}(x,Q^2) + g_2^{HT}(x,Q^2)\ , \tag{1.46}$$

where the first term proposed by Wandzura and Wilczek[36] contains only twist-2 matrix elements and is determined entirely by g_1 :

$$g_2^{WW}(x,Q^2) = -g_1(x,Q^2) + \int_0^1 \frac{dy}{y} g_1(y,Q^2). \tag{1.47}$$

The second term, g_2^{HT}, receives contributions from the quark-gluon interactions inside the nucleon and non-zero values of quark masses (twist-3 operators). In principle, twist-3 contributions can be big because of the strong confining interaction and the appreciable value of the s-quark mass. Bag model calculations support such an assumption.[35] If so, experimental measurements of g_2 directly measure the interaction-dependent higher twist matrix elements and could provide us with important information about nucleon structure.

1.3 QCD-improved parton model

1.3.1 Spin structure functions in QCD

In the "naive" parton model, the nucleon spin structure function g_1 defined by Eq. (1.30) is a weighted sum of the quark helicity distributions that depend only on the scaling variable x. In the QCD-improved parton model, quark distributions, and hence g_1 evolve with Q^2 due to gluon bremsstrahlung and gluon-induced quark-antiquark pair creation. In the leading order (LO) in the strong coupling constant α_S Eq. (1.30) simply becomes

$$g_1(x, Q^2) = \frac{1}{2} \sum_q e_q^2 \Delta q(x, Q^2). \tag{1.48}$$

The spin structure function g_1 is again defined by the quark contributions, and gluons couple to it only indirectly via the evolution of the quark distributions. In higher orders, g_1 is defined as

$$g_1(x, Q^2) = \frac{1}{2} \sum_q^{N_f} e_q^2 \left[C_q \otimes \Delta q + \frac{1}{N_f} C_G \otimes \Delta G \right] , \tag{1.49}$$

where again both quark and antiquark contributions are included into definition of $\Delta q \equiv \delta q + \delta \bar{q}$. The sum is over all active flavors ($N_f = 3$ is only light and strange quarks are taken into account), and the symbol $\otimes$ in Eq. (1.49) denotes the convolution integral

$$(f \otimes g)(x) = \int_x^1 \frac{dy}{y} f(y) g(\frac{x}{y}). \tag{1.50}$$

The perturbative Wilson coefficients $C_{q,G}$ represent QCD radiative corrections, and are written as series in powers of α_S

$$C_{q,G} = C_{q,G}^{(0)} + \frac{\alpha_S}{2\pi} C_{q,G}^{(1)} + \dots . \tag{1.51}$$

The leading order expression Eq. (1.48) is obtained by dropping higher-order $C_{q,G}^{(1)}$ terms and having $C_q^{(0)}(x) = \delta(1-x)$ and $C_q^{(0)}(x) = 0$. The next-to-leading (NLO) spin-dependent coefficients $C_q^{(1)}$ and $C_G^{(1)}$ in the modified minimal subtraction ($\overline{\mathrm{MS}}$) renormalization scheme are given[37] by

$$\begin{aligned} C_q^{(1)}(x) \quad &= \quad C_F \Bigg[(1+x^2) \left(\frac{\ln(1-x)}{1-x} \right)_+ - \frac{3}{2} \frac{1}{(1-x)_+} - \frac{1+x^2}{1-x} \ln x + \\ &\quad + 2 + x - \left(\frac{9}{2} + \frac{\pi^2}{3} \right) \delta(1-x) \Bigg] \end{aligned} \tag{1.52}$$

$$\text{and} \quad C_G^{(1)}(x) \;=\; 2T_f\left[(2x-1)\left(\ln\frac{1-x}{x}-1\right)+2(1-x)\right] , \tag{1.53}$$

and their moments are given in Appendix B. Here $C_F = 4/3$ and $T_f = N_f/2$ are Casimir invariants for the quark representation of $SU(3)_{flavor}$, and $(1-z)_+$ is defined by

$$\int_0^1 dz \frac{f(z)}{(1-z)_+} \equiv \int_0^1 dz \frac{f(z)-f(1)}{(1-z)}. \tag{1.54}$$

The charm contribution is usually ignored, so the number of flavors $N_f = 3$.

As it follows from Eq. (1.49), in the NLO analysis, the gluon density directly affects the spin structure function g_1. Precision measurements of g_1 can therefore reveal information about the gluon contribution to the nucleon helicity.

1.3.2 Q^2 dependence of the polarized parton distributions

In the simple "naive" parton model with no gauge degrees of freedom, the quark distributions are functions of only one variable – Bjorken x. This is not the case in QCD which predicts systematic deviations from this simple picture.

There are several sources of scaling violations. At high values of Q^2 quarks are likely to radiate gluons (Fig. 1.2[5]), and gluons, in turn, create quark-antiquark pairs. The gluon emission reduces the momentum of the quark effectively shifting the quark distribution to lower values of x. At high x the structure functions should then decrease with increasing Q^2, whereas at low x they are expected to increase. Thus, QCD predicts the *evolution* of the parton distributions with Q^2 that produces the logarithmic Q^2 dependence of the structure functions.

Another source of the scaling breaking is the effect of "higher twist" contributions, or corrections that arise from the subprocesses which involve more than the minimal number of interacting fields, such as the processes of Fig. 1.3. Although

---

[5]The figure shows virtual Compton scattering since its amplitude is related to the deep inelastic cross section via the optical theorem.



Fig. 1.2. Gluon emission results in logarithmic deviations from Bjorken scaling, as described by the DGLAP equations.



Fig. 1.3. Examples of higher twist corrections to the virtual Compton scattering amplitude.

these corrections are suppressed by powers of Q^2, they turn out to be significant at fixed $(1-x)Q^2$.[40]

1.3.2.1 Evolution of the parton distributions

Having the initial distributions at some value of $Q^2 = Q_0^2$ as the boundary conditions, one can obtain the distributions at higher Q^2 using the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) evolution equations[41–43] that read:

$$\begin{aligned} \frac{d\delta q_i(x,Q^2)}{d\ln Q^2} &= \frac{\alpha_S(Q^2)}{2\pi}\left[\delta q_i \otimes \Delta P_{qq} + \Delta G \otimes \Delta P_{qG}\right] \quad \text{and} \\ \frac{d\Delta G(x,Q^2)}{d\ln Q^2} &= \frac{\alpha_S(Q^2)}{2\pi}\left[\sum_{i=1}^{2N_f} \delta q_i \otimes \Delta P_{Gq} + \Delta G \otimes \Delta P_{GG}\right] \ , \end{aligned} \tag{1.55}$$

where (unlike the definition of Δ in Eq. (1.31)), $\delta q_i = q_i^{\uparrow}(x,Q^2) - q_i^{\downarrow}(x,Q^2)$ is the spin distribution for a quark (antiquark), $\Delta G = G^{\uparrow}(x,Q^2) - G^{\downarrow}(x,Q^2)$ is the gluon distribution, N_f is the number of active quark flavors, and $\otimes$ denotes the convolution integral of Eq. (1.50).

Like the perturbative Wilson coefficients of Eq. (1.51), the splitting functions are calculated as a series in the strong coupling constant

$$\Delta P_{ij} = \Delta P_{ij}^{(0)} + \frac{\alpha_S}{2\pi}\Delta P_{ij}^{(1)} + \ldots \, . \tag{1.56}$$

The leading order spin-dependent splitting functions are[41,43]

$$\begin{aligned}
\Delta P_{qq}^{(0)}(z) &= \frac{4}{3}\left[\frac{1+z^2}{(1-z)_+} + \frac{3}{2}\delta(z-1)\right] \, , \\
\Delta P_{qG}^{(0)}(z) &= \frac{1}{2}\left[z^2 - (1-z)^2\right] \, , \\
\Delta P_{Gq}^{(0)}(z) &= \frac{4}{3}\left[\frac{1-(1-z)^2}{z}\right] \, , \quad \text{and} \\
\Delta P_{GG}^{(0)}(z) &= 3\left[(1+z^4)\left(\frac{1}{z} + \frac{1}{(1-z)_+}\right) - \frac{(1-z)^3}{z} + \left(\frac{11}{6} - \frac{N_f}{9}\right)\delta(1-z)\right] \, .
\end{aligned} \tag{1.57}$$

The next-to-leading order spin-dependent splitting functions have been calculated in Ref. [37], and their moments are given in Appendix B. The evolution equations for the unpolarized distributions can be easily obtained by changing $\delta q_i \to q_i$, $\Delta G \to G$, $\Delta P_{ij} \to P_{ij}$, and the unpolarized splitting functions $P_{ij}^{(0)}$ given in References [41,43] (LO), and References [38,39] (NLO).

It is not possible to find an exact analytic solution to the Equations (1.55). However, numerical calculations and approximations are feasible.[44] It is clear from Eqs. (1.55) that the evolution of the quark distributions is sensitive to the gluon polarization; thus studying the Q^2 dependence of the polarized structure functions can provide us with the experimental constraints on the polarized gluon distribution functions. For the presently available experimental data, such a program is carried out in Chapter 5.

It should be noted that in the leading order, qq elements of the evolution matrix are equal in both polarized and unpolarized cases: $\Delta P_{qq}^{(0)} = P_{qq}^{(0)}$. Thus, if the gluon contribution is small (such is the case at high x), the evolution of the unpolarized and polarized structure functions is similar, and the ratio g_1/F_1 (or the asymmetry A_1) is approximately independent of Q^2. This observation, supported within the precision of the data, has been used by the experimental collaborations to evolve data taken at different values of Q^2 to a common scale. This assumption is clearly unjustified at low x where the gluon densities, both unpolarized and polarized, rise sharply. As the quality of the data improves, the Q^2-coverage increases, and even lower values of x are probed, a consistent pQCD-based approach should be used.

1.3.2.2 Higher twist effects

The diagrams involving interference terms between different quark currents as well as the quark-gluon correlations (Fig. 1.3) are responsible for the contributions to the structure functions that are suppressed relative to the leading contributions by the powers of Q^2. These corrections, however, become important at fixed values of W^2. Taking the *higher twist* corrections into account leads to the following series expression for the unpolarized structure functions[40]

$$F_2(x, Q^2) = F_2(x, Q^2)_{\text{leading twist}} \left[1 + \frac{A}{Q^2(1-x)} + \frac{B}{Q^4(1-x)^2} + \cdots \right] , \qquad (1.58)$$

where the coefficient $A \approx O(m_\rho^2)$ is set by the wavefunction scale. Unpolarized data from SLAC show substantial higher twist contributions at high x.[79] One can therefore assume that higher twist contributions to the polarized structure functions are not negligible at low W^2.

Higher twist corrections modify the structure function sum rules as follows:

$$\Gamma(Q^2) = \Gamma_\infty \left[1 + \sum_n C_n \left(\frac{\alpha_S(Q^2)}{\pi} \right)^n \right] + \sum_m \frac{D_m}{(Q^2)^m} , \qquad (1.59)$$

where D_m are the higher twist coefficients. There is no firm agreement on the size of the correction for the Bjorken sum rule; the recent estimate[78] is

$$D_1 = -0.02 \pm 0.01 \text{ GeV}^2 \tag{1.60}$$

so the effect is quite small but comparable to the size of the current experimental errors.

1.4 Shape of the polarized parton distributions

1.4.1 Helicity-dependent quark distributions

The Q^2-evolution of structure functions is well predicted by perturbative QCD DGLAP equations.[41–43] The initial shape of these distributions, however, is not directly calculable and reflects the non-perturbative dynamics of quarks and gluons inside the nucleon. Nevertheless, the end-point behavior of the structure functions can be accurately predicted from perturbative arguments. In this section, we will constrain the behavior of the structure functions in the regions of $x \approx 0$ and $x \approx 1$. The predicted forms can be combined to form a parameterization of the polarized structure functions.

The polarized quark and gluon distribution functions $G_{q/H}(x, \lambda, Q)$ and $G_{g/H}(x, \lambda, Q)$ of a hadron are most conveniently represented as overlaps of the light-cone wave functions $\psi_n(x_i, k_\perp, \lambda_i)$, where[6] $\sum_{i=1}^{n} x_i = 1$, and $\sum_{i=1}^{n} k_\perp = 0$. As shown in Appendix A, the region of $x \to 1$ represents a very far off-shell configuration of the bound state wave function. Assuming that the bound state wave function is dominated by the lowest invariant mass Fock state, one can conclude that a constituent can get $x \approx 1$ only by exchanging hard gluons. Thus, the leading behavior of the amplitudes in the region of $x \approx 1$ can be computed from the simple minimally connected gluon-exchange diagrams (Fig. 1.4) calculable in perturbative QCD.

[6]see Appendix A



Fig. 1.4. Leading-order diagram for $G_{q/H}(x, \lambda, Q)$ at $x \approx 1$.

The limiting power-law behavior at $x \to 1$ of helicity-dependent distributions is

$$G_{q/H} \sim (1-x)^{2n-1+2\Delta S_z}. \tag{1.61}$$

Here n is the minimal number of spectator quark lines, and $\Delta S_z = \mid S_z^q - S_z^H \mid = 0, 1$ for parallel or anti-parallel quark and proton helicities, respectively.[50] This counting rule (*cf.* Eqs. (A.13), (A.14), and (A.15)) reflects the fact that the valence Fock states with the minimum number of constituents give the leading contribution to structure functions when one quark carries nearly all of the light-cone momentum; contributions from Fock states with a higher number of partons fall off faster at $x \to 1$.[49] The helicity dependence of the counting rule also reflects the helicity retention properties of the gauge couplings; a quark with a large momentum fraction of the hadron also tends to carry its helicity. The anti-aligned helicity quark is suppressed by a relative factor $(1-x)^2$. Similarly, in the case of splitting functions such as $q \to qg$ or $g \to \bar{q}q$, the sign of the helicity of the parent parton is transferred to the constituent with the largest momentum fraction.[51] The counting rule for valence quarks can be combined with the splitting functions to predict the $x \to 1$ behavior of gluon and non-valence quark distributions of non-exotic hadrons; they fall off faster by at least one power of $(1-x)$ than the respective quark distributions.

The counting rules for the end-point-behavior of quark and gluon helicity distributions can also be derived from duality, *i.e.* a direct relationship between the physics of exclusive and inclusive channels at fixed invariant mass.[52] As shown by Drell and Yan,[52] a quark distribution function behaves as

$$G_{q/H} \sim (1-x)^{2n-1} \tag{1.62}$$

at $x \to 1$ if the corresponding form factor falls off as

$$F(Q^2) \sim (1/Q^2)^n \tag{1.63}$$

at large Q^2. Measurements of elastic electron-proton scattering at SLAC[53] are compatible with the perturbative QCD predictions[54] for both the helicity-conserving $F_1(Q^2)$ and helicity-changing $F_2(Q^2)$ form factors: $Q^4 F_1(Q^2)$ and $Q^6 F_2(Q^2)$ become approximately constant at large Q^2. This behavior corresponds to the threshold behavior of helicity-parallel and helicity anti-parallel quark distributions: $(1-x)^3$ and $(1-x)^5$ as $x \to 1$, respectively, in agreement with the counting rules. The leading exponent for quark distributions is odd in the case of baryons and even for mesons in agreement with the Gribov-Lipatov crossing rule.[41]

The counting rule predictions for the quark and gluon distributions are relevant at low momentum transfer scales $Q_0 \approx \Lambda_{QCD}$ in which the controlling physics is that of the hadronic bound state rather than the radiative corrections associated with structure function evolution.

At high Q^2 the radiation from the struck quark line increases the effective power law fall-off $(1-x)^p$ of structure functions relative to the underlying quark distributions[49]:

$$\Delta p = (4C_F/\beta_1) \ln \left(\frac{\ln(Q^2/\Lambda^2)}{\ln(Q_0^2/\Lambda^2)} \right) , \tag{1.64}$$

where $C_F = 4/3$ and $\beta_1 = 11 - (2/3)N_f$. Thus, the counting rule predictions for the power p provide a *lower bound* for the effective exponent of quark structure functions at high $Q^2 > Q_0^2$. However, in the end-point region $x \approx 1$, the struck quark is far off-shell and the radiation is quenched since one cannot evolve Q^2 below

$Q_0^2 \simeq k_F^2 = -(\mu^2/(1-x))$, the Feynman virtuality of the struck parton[40] (here, μ is the invariant mass of the spectator quarks).

Thus pQCD can give useful predictions for the power law fall-off of helicity-aligned and anti-aligned structure functions at $x \approx 1$. Higher order contributions involving additional hard gluon exchange are suppressed by powers of $\alpha_S(k_F^2)$. Further iterations of the interaction kernel will give factors of fractional powers of $\log(1-x)$ analogous to the anomalous dimensions $\log^{\lambda_n} Q^2$ which appear in the pQCD treatment of form factors at large momentum transfer.[54]

The fact that one has a definite prediction for the $x \approx 1$ behavior of leading twist structure functions is a powerful tool in QCD phenomenology, since any contribution that does not decrease sufficiently fast at large x is most likely due to coherent multi-quark and quark-gluon correlations. As discussed in Ref. [55], such contributions are higher twist, and they are significant at fixed $(1-x)Q^2 \approx M^2$.

1.4.2 Helicity-dependent gluon distributions

At $x \to 1$ the gluon distribution function is constrained by the counting rule arguments (see Fig. 1.5) and obeys the Gribov-Lipatov crossing rule[41]:

$$G_{g/p} \sim (1-x)^4, \quad x \to 1. \tag{1.65}$$

It is known in QED[51] that in $e \to e\gamma^*$ processes the polarization of the initial electron is retained by the photon at high momentum transfer; the same behavior is expected for any gauge theory including QCD. Thus,

$$\frac{G^-(x)}{G^+(x)} = (1-x)^2, \quad x \to 1. \tag{1.66}$$

At low x quarks radiate coherently (Fig. 1.6), and one has to take into account interference terms. Brodsky, Burkardt, and Schmidt have obtained the following constraint on the gluon distribution functions at $x \approx 0$[49]:

$$\left(\frac{\Delta G(x)}{G(x)}\right)_{\text{proton}} \to \frac{x}{3}\left\langle\frac{1}{y}\right\rangle, \quad x \to 0. \tag{1.67}$$



Fig. 1.5. The leading contribution to the gluon distribution function at high x.



Fig. 1.6. The low-x behavior of the gluon distribution function is determined by coherence effects.

Here $\langle 1/y \rangle$ stands for the first inverse moment of the quark light-cone momentum fraction distribution in the lowest proton Fock state. For this state $\langle 1/y \rangle \approx 3$.

For baryons, a simple form of the gluon distributions, which incorporates the limiting behaviors presented above, is[49]

$$\Delta G(x) = \frac{N}{x}[1 - (1-x)^2](1-x)^4, \tag{1.68}$$

$$G(x) = \frac{N}{x}[1 + (1-x)^2](1-x)^4 \ , \tag{1.69}$$

where N is a normalization constant. In this model the momentum fraction carried by the gluons in the proton is

$$\langle x_g \rangle \equiv \int_0^1 dx \ xG(x) = \frac{12}{35}N, \tag{1.70}$$

and the helicity carried by the gluons is

$$\Delta G \equiv \int_0^1 dx \Delta G(x) = \frac{11}{30} N, \tag{1.71}$$

Taking the momentum fraction $\langle x_g \rangle$ to be 1/2 yields $\Delta G = 0.54$.

1.4.3 The axial anomaly

The prediction, found in the previous section, that $\Delta G \simeq 0.5$ is phenomenologically interesting. If one also accepts the experimental suggestion from EMC that the quark helicity sum $\Delta u + \Delta d$ is small, then this implies that gluons could carry a large part of the proton helicity $J_z = 1/2$. The angular momentum conservation requires then significant (and, most likely, negative) orbital angular momentum L_z which arises, for example, from the finite transverse momentum associated with the $q \to qg$ gluon emission matrix element.

Several authors have argued[56–58] that the gluon polarization may explain the small quark helicity content of the proton observed by EMC and rescue the Ellis-Jaffe assumption of $\Delta s = 0$.[59] Gluons could contribute to proton helicity via the γ_5-triangle anomaly that breaks, at the quantum level, the conservation of the axial current $\bar{q}\gamma_\mu\gamma_5 q$:

$$\partial^\mu A^0_\mu = \frac{\alpha_s}{\pi} \; N_f \; \mathrm{tr} \; F_{\mu\nu}\tilde{F}^{\mu\nu} \; , \quad \tilde{F}^{\mu\nu} \equiv \epsilon^{\mu\nu\alpha\beta} F_{\alpha\beta} \tag{1.72}$$

where $F_{\mu\nu}$ is the gauge field strength tensor in QCD. The intuitive meaning of Eq. (1.72) is that the anomaly induces a mixing between gluons and the flavor-singlet axial current of quarks. One can then redefine the quark helicities measured in the experiment to be

$$\Delta q \to \Delta q - \frac{\alpha_S}{2\pi} \Delta G. \tag{1.73}$$

The authors of Ref. [60] have calculated, based on the recent experimental data,that the $\Delta s = 0$ assumption requires $\Delta G \approx 2$. This number is not as big as claimed in Ref. [59], but still requires a large negative orbital angular momentum contribution.

Equation (1.73) demonstrates the intrinsic ambiguity associated with the interpretation of the quark distribution function. This is the manifestation of the *factorization scheme dependence* of the parton distributions.[61–63] Although the structure functions measured in the experiment are unambiguous, their factorization in Eq. (1.49) into non-perturbative parton distributions Δf_i and hard cross sections ΔC_i is definition-dependent. One therefore always has to specify a particular prescription (*scheme*) according to which the factorization is performed. In fact, if the calculation is done using the dimensional regularization in the conventional $\overline{\mathrm{MS}}$ scheme, the gluon contribution to the first moment of g_1 vanishes[61] (due to the fact that $\int_0^1 dx C_G(x) = 0$ in Eq. (1.49)[37]). We will discuss more details of the scheme dependence as well as other theoretical uncertainties in extracting the parton distributions in Chapter 5.

1.5 Low and high x phenomenology

No experiment can measure the full range $0 \le x \le 1$, and every experiment must make some assumption in order to extrapolate to the full x range. The contribution from the high x is typically small since the structure functions decrease rapidly. From the perturbative QCD arguments outlined in Section 1.4.1, the valence quark contribution to g_1 falls off at least as $(1-x)^3$ at $x \to 1$, and the sea and gluon contributions are suppressed by additional powers of $(1-x)$. The helicity-antialigned quark distribution is suppressed by $(1-x)^2$ relative to the helicity-aligned distribution, so one can write on general grounds for the virtual photon-nucleon asymmetry

$$A_1 \sim \frac{\Delta q}{q} \to 1, \; x \to 1. \tag{1.74}$$

Experiments commonly use Eq. (1.74) to extrapolate data to $x = 1$, or extrapolate g_1 directly assuming the pQCD predicted power fall off

$$g_1 \sim \Delta q \sim (1-x)^3, \; x \to 1. \tag{1.75}$$

Both extrapolations are consistent with the data within experimental errors.

The situation is much less clear for the small x extrapolation. At fixed Q^2, the $x \to 0$ limit corresponds to the infinite photon energy $\nu \to \infty$, or, equivalently,

$$s \equiv (P+q)^2 \equiv W^2 = M^2 - Q^2 + 2M\nu = M^2 + Q^2 \frac{1-x}{x} \to \infty \tag{1.76}$$

for the photon-nucleon center-of-mass energy s. This limit has been traditionally described by the theory of complex angular momentum (Regge theory). In Regge theory, one assumes that the high energy limit of the s-channel amplitude for the scattering $ab \to cd$ is determined by the singularities of the t-channel amplitude in the complex angular momentum plane; in the case of a simple pole (associated with an exchange of a particle – *"Reggeon"*[7]), the corresponding contribution to the amplitude can be written as

$$\mathcal{A}^{ab\to cd}(s,t) = \beta(t) s^{\alpha(t)}, \; s \to \infty \;, \tag{1.77}$$

where $\alpha(t)$ is the Regge *trajectory.* The optical theorem relates the total cross section to the imaginary part of the forward ($t=0$) amplitude:

$$Im\mathcal{A}^{ab\to cd}(s,t=0) = s\sigma_{tot}^{ab\to cd}(s), \tag{1.78}$$

and we have

$$\sigma_{tot}^{ab\to cd}(s) \sim s^{\alpha(0)-1}. \tag{1.79}$$

The Regge *intercept* $\alpha(0)$ is determined experimentally, for instance, in πN or NN scattering.

Regge theory has been very successful in describing a wide variety of high energy *unpolarized* cross sections.[64] Since it provides a simple and economical (in terms of free parameters) description of the total cross sections, Donnachie and Landshoff conclude that[64] *"Regge theory remains one of the great truths of particle physics".*

[7]Such a particle can possess quantum numbers of a real meson, or vacuum, in which case it is called a Pomeron[32].

In Section 1.2.2 we have shown the relationship between the virtual nucleon photoabsorption cross sections and the spin structure functions. For $\nu \to \infty$ and a fixed Q^2, the leading expression for g_1 is

$$g_1 \approx \nu(\sigma^T_{1/2} - \sigma^T_{3/2})/2 \tag{1.80}$$

(ignoring terms constant or decreasing with ν). Since $s \approx 2M\nu$ at large ν and fixed Q^2,

$$g_1 \sim \beta(Q^2)\nu^\alpha \sim x^{-\alpha},\ x \to 0\ , \tag{1.81}$$

where α is the dominant Regge intercept and $\beta(Q^2)$ is the pole residue. It is believed[65] that the leading behavior of g_1 is controlled by the a_1 meson trajectory with the intercept $\alpha_{a_1} = -0.14 \pm 0.20$.[65,66] Thus g_1 converges at low x according to the traditional Regge picture. Experiments so far have used the upper limit $\alpha_{a_1} = 0$ to extrapolate data to $x = 0$ and evaluate sum rules; the variation in α_{a_1} is used to estimate the extrapolation error.

There are several problems with this simple description. It is not certain at which values of x, and at what Q^2 the Regge theory is applicable. Strictly speaking, it works at the low hadron scales, *i.e.* Q^2 below 1 GeV^2. At high ν and Q^2 and finite x the structure functions are, in general, not determined by the rightmost singularities in the complex angular momentum plane; one has to include all relevant contributions. One therefore has to assume[32] that there is a smooth transition from the scaling regime ($\nu \to \infty$, $Q^2 \to \infty$, x fixed) to the Regge regime ($\nu \to \infty$, $Q^2 = const$) at small x. This is an additional hypothesis. In the unpolarized case, it has been shown[32] that the residues of the dominating Regge poles $\beta(Q^2)$ have to have a special Q^2 dependence. Even if the Regge regime is not truly set in, the unpolarized analysis benefits from the fact that the residues of different poles are generally of the same sign (positive), whereas in the polarized case this may not be the case and one cannot ignore possible "interference effects" that could mask the true asymptotic behavior.

Several authors argued[67,68] that the double-Pomeron cut contributes to g_1 at low x; it then induces a singular behavior[8]

$$g_1 \sim \frac{1}{x \log^2 x}, \; x \to 0. \tag{1.82}$$

Again, the kinematic domain where such behavior applies is not specified; moreover, it has not been observed in the unpolarized data.[69]

A model of the Pomeron based on the non-perturbative gluon exchange[70] gives a softer, but still singular behavior:

$$g_1 \sim 1 + 2 \log x, \; x \to 0. \tag{1.83}$$

The coherence arguments based on perturbative QCD and outlined in Section 1.4.2 imply that at some low hadron scale the polarized gluon distribution $\Delta G(x)$ is related to the unpolarized one $G(x)$ according to the Eq. (1.67), *i.e.*

$$\Delta G(x) \sim x G(x), \; x \to 0. \tag{1.84}$$

Since the evolution mixes quark singlet and gluon distributions (see Section 1.3.2.1), the most singular of the two controls the behavior of the structure functions at low x and moderately high Q^2. Thus, if the behavior of $G(x)$ is harder than $1/x$, $\Delta G(x)$ and consequently $g_1(x)$ is singular at low x.

Finally, instead of relying in the theoretical predictions, one can fit the data to a phenomenological model. A power-law dependence similar to Eq. (1.81) but with power α varying freely is a reasonable choice. On general grounds, one should distinguish between isospin 1 and isospin 0 states (or, in QCD language, between non-singlet and singlet contributions to g_1). In this case, the fit of the following form is performed:

$$g_1 = b_{NS} x^{-a_{NS}} + b_S x^{-a_S} \, , \tag{1.85}$$

where S stands for singlet and NS for non-singlet quark distributions.

[8] There are also counter-arguments, see, for example Ref. [32] and Ref. [66].

There exists a theoretical estimate[71] of the powers a_{NS} and a_S based on the summation of the contributions of the form $\alpha_S \log^2(1/x)$. It was shown to produce a very divergent behavior

$$g_1^{NS} \sim \frac{1}{x^{0.4}},\ g_1^S \sim \frac{1}{x^{1.01}\log^{3/2}(1/x)},\ x \to 0. \tag{1.86}$$

This is much more singular than the dependence induced by the GLAP evolution.

Clearly, precise experimental data at high energies are needed to determine the low x behavior of the polarized structure functions. So far the data have not been able to constrain it well, which poses certain problems in evaluating the polarized sum rules. We will return to this issue in Section 4.4.3.

1.6 Extracting g_1^n from ^{3}He data

In the experiment E154, we scattered polarized electrons off a polarized ^{3}He target. To a very good approximation, ^{3}He nucleons are in the spatially symmetric S state. The two protons are in a symmetric isospin state, and therefore, due to the Pauli principle, they are in the antisymmetric spin state. The total spin of ^{3}He is thus carried by the neutron in this approximation, and asymmetries measured on a ^{3}He target are directly proportional to the neutron structure functions.

In a more realistic model, other components of the ^{3}He wave function are included. Besides the S-wave, the three-body wavefunction contains percentages of the S' wave, arising from the differences in the tensor $T = 0$ and $T = 1$ forces distorting the primary S state, and the D wave.[72] The P-wave is suppressed due to its opposite parity. The admixture of S' and D partial waves reduces the net neutron polarization p_n in ^{3}He and introduces finite proton polarization p_p. The proton and neutron polarizations can be calculated by considering the quantities $P_{p(n)}^{(\pm)}$, representing the probability to have a proton (neutron) with spin parallel (+) or antiparallel (−) to the ^{3}He spin. In a pure S wave state $P_n^{(+)} = 1$, $P_n^{(-)} = 0$ and

$P_p^{(+)} = P_p^{(-)} = \frac{1}{2}$, whereas for a three-body wave function containing S, S' and D waves, one has[72,73]

$$\begin{aligned} P_n^{(+)} &= 1 - \Delta \\ P_n^{(-)} &= \Delta \\ P_p^{(\pm)} &= \frac{1}{2} \mp \Delta', \end{aligned} \tag{1.87}$$

where $\Delta = \frac{1}{3}[P_{S'} + 2P_D]$ and $\Delta' = \frac{1}{6}[P_D - P_{S'}]$. Here we introduced $P_{S'}$ and P_D as probabilities of S' and D state, respectively.

From published calculations on the three body system one obtains, in correspondence of the experimental value of the binding energy of ^{3}He, $\Delta = 0.07 \pm 0.01$ and $\Delta' = 0.014 \pm 0.002$.[72] In the approximation of *independent scattering* off unbound nucleons, one can write:

$$g_{1^3\mathrm{He}}(x) = 2p_p g_1^p(x) + p_n g_1^n(x) \tag{1.88}$$

and

$$A_{^3\mathrm{He}} = (2f_p p_p A_p + f_n p_n A_n) \ , \tag{1.89}$$

where A is the measured asymmetry $A_{\|}$ or $A_{\perp}$, and $f_{p(n)} = F_2^{p(n)}/[(2F_2^p + F_2^n) f_{\mathrm{EMC}}]$ is the proton (neutron) dilution factor. The nuclear effects in the unpolarized scattering are taken into account by the EMC effect factor.[74] Taking into account the S' and D state probabilities, the effective nucleon polarizations are:

$$\begin{aligned} p_p &= P_p^{(+)} - P_p^{(-)} = -0.028 \pm 0.004 \quad \text{and} \\ p_n &= P_n^{(+)} - P_n^{(-)} = 0.87 \pm 0.02. \end{aligned} \tag{1.90}$$

In the assumption of independent scattering, one neglects nuclear binding effects and effects of Fermi motion and nuclear shadowing. It was shown by the authors of Ref. [73] that for the polarization observables these effects are only noticeable for $x > 0.9$ and $x \approx 0$ in the deep inelastic scattering regime $Q^2 > 1$. We use Eq. (1.89) to extract neutron asymmetries from the ^{3}He data. Thus, polarized ^{3}He is good model of polarized neutron.

1.7 Experimental status

The deep inelastic scattering program at SLAC was started in the late 1960's with the SLAC-MIT experiment that led to the discovery of the quark-parton model.[2–5] The first experiment on spin-dependent deep inelastic scattering was E80 at SLAC in 1976.[6] The experiment used a polarized electron beam with energies of 9.7 and 12.9 GeV and a polarized butanol target to measure the photon-nucleon asymmetry A_1^p. The technique of Dynamical Nuclear Polarization[75] was used to polarize protons to $\approx$ 50%. The experiment was repeated in 1980 by the same SLAC-Yale collaboration (SLAC E130)[7] with higher electron energies (16.2 and 22.7 GeV). Both experiments used fixed-angle magnetic spectrometers to detect scattered electrons, centered at 9° (E80) and 10° (E130). They found good agreement with the quark-parton expectations over the covered x range. The main parameters of E80 and E130, as well as of other experiments to be described in this Section, are summarized in Table 1.2.

After the early series of the experiments on the spin-dependent deep inelastic scattering ended at SLAC in early 1980's, the experimental program was continued at CERN. The EMC experiment took data in 1984/5, and the first results became available in 1988.[8] EMC scattered a μ^+ beam with energies between 120 and 200 GeV off a polarized ammonia target. The muon beam is naturally polarized to $\approx$ 80% since it is produced in the π^+ decay. The high muon energy allowed the measurement to extend to much lower x than in original SLAC experiment, *i.e.* values of $x \approx 0.01$. The EMC results was in good agreement with the naive quark model expectations and SLAC results for $x > 0.1$, but the data lay significantly below the prediction at lower values of x. Consequently, combining EMC and SLAC

Table 1.2. Parameters of polarized deep-inelastic scattering experiments.

Experiment	years	target	E_b (GeV)	P_b	P_t	Events (10^6)
SLAC E80	1976	butanol	9 – 12	50%	50% (p)	2
SLAC E130	1980	butanol	16 – 22	80%	60% (p)	1
CERN EMC	1988	ammonia	190	80%	75%	1
SLAC E142	1992	^{3}He	23	35%	33%	300
SLAC E143	1993	ammonia (p)	29	80%	80%	200
		ammonia (d)			40%	
CERN SMC	1992–	butanol (d)	190	80%	40%	≈20
	1996	ammonia (p)			80%	
DESY HERMES	1995–	atomic	27	60%	40% (^{3}He)	≈6
		^{3}He, ^{1}H, ^{2}D				
SLAC E154	1995	^{3}He	48	80%	38%	100
SLAC E155	1997	ammonia (p)	48	80%	80%	???
		^{6}LiD (d)			50%	

data and using a smooth Regge extrapolation $g_1^p \sim x^{0.12}$, $x \to 0$, EMC obtained for the first moment of g_1^p

$$\Gamma_1^p(\mathrm{EMC}) = 0.126 \pm 0.010 \text{ (stat.) } \pm 0.015 \text{ (syst.)} \tag{1.91}$$

at an average $Q^2 = 10$ GeV^2, about a three standard deviation disagreement with the Ellis-Jaffe prediction. Assuming $SU(3)_{\mathbf{flavor}}$ symmetry and using Eq. (1.42), one can find that the total quark contribution to the proton spin is small:

$$\Delta\Sigma = 0.12 \pm 0.17. \tag{1.92}$$

This result came as a surprise, and the effect was even dubbed "the proton spin crisis" in the community. It has inspired a large amount of theoretical work aimed at understanding the spin structure of the proton. It has also been the genesis of a new experimental program in polarized DIS. The interest shifted towards experimental tests of the Bjorken sum rule, and precision determinations of the spin structure of the nucleon.

The SLAC spin structure program restarted in the Fall of 1992 with the experiment E142.[9] The experiment used a AlGaAs polarized electron source developed for the SLC which produced a high-intensity beam of 40% polarization. The beam was delivered to the End Station A with energies of 19, 22, and 25 GeV. A high-density polarized ^{3}He target was developed for this experiment, for which ^{3}He gas was polarized using optical pumping and spin exchange techniques (see Section 2.5) to yield an average polarization of 33%. Scattered electrons were detected in two independent magnetic spectrometers centered at 4.5° and 7°.[76] The experiment collected approximately 300 million deep-inelastic events over the kinematic range $0.03 \leq x \leq 0.6$ at the average Q^2 of 2 GeV^2, which resulted in the most precise determination of the neutron structure function g_1^n, prior to E154. Unlike the original SLAC experiments, the transverse asymmetry $A_\perp$ was also measured to minimize the systematic error in extracting g_1^n. An additional advantage of the SLAC experiments was the ability to flip the electron helicity pseudo-randomly on the pulse-by-pulse basis. Combined with the regular reversals of the target spin orientation, this feature resulted in the strong suppression of the beam-related systematic effects.

Experiment E143 at SLAC ran in the winter of 1993/4.[10,11] The development of strained $GaAs$ photocathodes resulted in high beam polarization (80%) available to E143. The beam energy was raised to 29 GeV; data were also taken at lower beam energies of 9 and 16 GeV to study the Q^2 dependence of the spin-dependent structure function g_1.[77] Polarized ammonia and deuterated ammonia targets were used to measure the spin dependent structure functions of the proton and deuteron. The spectrometer setup was essentially unchanged from E142. The experiment collected approximately 200 million deep inelastic events, and both the longitudinal[10,11] and the transverse[22] structure functions were measured over the range $0.03 \leq x \leq 0.7$ at an average Q^2 of 3 GeV^2.

The program at CERN continued with the SMC experiment[12,13] that took data with polarized deuteron targets in 1992, 1994, and 1995 (deuterated butanol was used as a target material), and with a proton target in 1993 and 1996. Like EMC, SMC used the highest energy 190 GeV muon beam, and the butanol target was superseded in 1996 by an ammonia target. Due to the high beam energy, the experiment reached lower values of x and higher Q^2 than the SLAC experiments. The measurements covered the x range of $0.003 \leq x \leq 0.7$ at an average Q^2 of 10 GeV^2. However, the muon intensity was low, and statistics limits the precision of the SMC measurements.

A new spin structure program was started recently at DESY. The HERMES experiment[80] operates in the HERA storage ring utilizing the 27 GeV positrons (electrons) and polarized internal gas targets. The novel technique is used to inject the polarized atoms into the windowless target chamber inside the storage ring, thus allowing to have the dilution factor very close to unity. The main feature

Table 1.3. Values of the first moments of g_1^p, g_1^n, and g_1^d reported by experiments at SLAC and CERN.

Experiment	Target	Q^2 (GeV^2)	Γ_1	Ref.
CERN EMC	p	10	0.126 ± 0.010 (stat.) ± 0.015 (syst.)	[8]
CERN SMC	p	10	0.136 ± 0.011 (stat.) ± 0.011 (syst.)	[12]
CERN SMC	d	10	0.041 ± 0.006 (stat.) ± 0.005 (syst.)	[13]
SLAC E142	n	2	0.031 ± 0.006 (stat.) ± 0.009 (syst.)	[9]
SLAC E143	p	3	0.129 ± 0.004 (stat.) ± 0.009 (syst.)	[10]
SLAC E143	d	3	0.042 ± 0.003 (stat.) ± 0.004 (syst.)	[11]

of the experiment is the ability to identify the final state hadrons simultaneously with scattered positrons. By tagging the flavor of the leading hadron (*i.e.* doing a *semi-inclusive* measurement), one is able to probe directly the valence and sea quark distributions inside the nucleon. The first inclusive results from the 1995 run with the ^{3}He target have been released in the preliminary form[81] and are expected to be published soon. The experiment will continue to run into the next century.

The summary of the data available prior to E154 is given in Table 1.3 and Figures 1.7 and 1.8. The consistency of the data taken in different experiments and at different kinematics, is outstanding. Not only the tests of sum rules are possible, but information about the shape of the structure functions, and the underlying parton distributions has begun to emerge.

Barring difficulties with the low-x extrapolation and the Q^2 evolution of the structure functions, the values of the moments in Table 1.3 can be used to test the Bjorken sum rule and extract the total quark contribution to the proton helicity $\Delta\Sigma$. Ellis and Karliner[82] performed a global fit to the data and obtained

$$\int_0^1 dx\,\left[g_1^p(x,Q^2) - g_1^n(x,Q^2)\right] = 0.164 \pm 0.011 \tag{1.93}$$

at $Q^2 = 3$ GeV^2, in perfect agreement with the prediction. They went even further, and assuming the validity of the Bjorken sum rule, extracted the value of α_S from the Q^2 dependence of the measured quantity (Eq. (1.41)):



Fig. 1.7. World data on $xg_1^p(x)$.

$$\alpha_S(3\ \mathrm{GeV}^2) = 0.328^{+0.026}_{-0.037}\ , \tag{1.94}$$

which is quite competitive with other determinations which are available.[20] For the quark helicity contributions, Ellis and Karliner obtained[82]

$$\begin{aligned} \Delta u &= 0.82 \pm 0.03 \\ \Delta d &= -0.44 \pm 0.03 \\ \Delta s &= -0.11 \pm 0.03 \end{aligned} \tag{1.95}$$

and

$$\Delta\Sigma = \Delta u + \Delta d + \Delta s = 0.27 \pm 0.04. \tag{1.96}$$

One has to keep in mind, of course, that the theoretical errors associated with the low x extrapolation, Q^2 dependence of asymmetries, higher twist effects, etc. have



Fig. 1.8. World data on $xg_1^n(x)$.

not been included in Equations (1.96) and (1.96).

Several pQCD fits to the polarized deep inelastic data have been made in the leading[60,83–85] and next-to-leading[63,86,87] orders in α_S. The first information on the polarized parton distributions have become available. We will discuss this program in Chapter 5.

Chapter 2

Experimental Apparatus

2.1 Overview

In SLAC experiment E154 the spin dependent structure functions $g_1^n(x, Q^2)$ and $g_2^n(x, Q^2)$ of the neutron were measured in the kinematic range $0.014 \leq x \leq 0.7$ and $1\ \mathrm{GeV}^2 \leq Q^2 \leq 17\ \mathrm{GeV}^2$ using a polarized, optically pumped ^{3}He gas target and a longitudinally polarized 48.3 GeV electron beam. The average target polarization of 38% was achieved in the longitudinal direction, and the average beam polarization was determined to be 82%. Scattered electrons were detected simultaneously in two independent large acceptance magnetic spectrometers centered around 2.75° and 5.5° relative to the beam line. Results for g_1^n were obtained by measuring the asymmetry between the cross sections in which the target and beam polarizations were parallel versus anti-parallel and using existing measurements of the unpolarized structure functions F_1 and F_2.

The experiment was a collective effort of 80 physicists from 22 institutions. It ran for two months in October-November, 1995. In this Chapter, we describe the apparatus used to perform the measurement.

2.2 Polarized electron source and beam transport

2.2.1 Polarized electron source

The polarized electrons were produced by photoemission from a strained GaAs photocathode.[88,89] The photocathode was illuminated by a flashlamp-pumped Ti:sapphire laser that produced $\approx 10\ \mu$s long pulses at a wavelength of 850 nm with a 120 Hz frequency. The pulses were then sliced by the Laser Pulse Chopper (LPC) to



Fig. 2.1. A schematic of the polarized source at SLAC. The setup shown is for the SLC operation. The E154 setup was very similar with one of the lasers changed to flashlamp-pumped Ti:sapphire.

200 – 250 ns and shaped slightly non-uniformly to compensate for the beam loading effects in the accelerator and to obtain uniform pulses in the end of the linac. A schematic of the polarized source is presented in Fig. 2.1.

The laser light was polarized with a linear polarizer and a combination of two Pockels cells, the quarter-wave plates that can change the optical axes depending on the applied high voltage. The axes of one of the Pockels cells were rotated by 45% relative to the axes of the linear polarizer and the other cell. This cell (CP Pockels cell) was used to generate the circularly polarized light; the sign of the high voltage determined the helicity of the light incident on the photocathode (and hence the helicity of the electrons). The second cell (PS Pockels cell) was used to control the charge asymmetry of the beam.



Fig. 2.2. Energy levels in strained GaAs.

The energy levels of a *strained* GaAs are illustrated in Fig. 2.2. The strain is induced by growing a 0.1 μm layer of GaAs on a substrate of GaAsP. The difference in the lattice constants of GaAs and GaAsP creates the compressive strain and splits the degeneracy of the $J = 3/2$ valence band by $\approx$ 50 meV. Photons with positive helicity and energy 1.43 eV $< E_\gamma <$ 1.48 eV excite the transition from the $m_j = -3/2$ valence band level to the $m_j = -1/2$ conduction band level. The electrons then diffuse from the conduction level into the vacuum. The quantum efficiency (QE) of the cathode is increased by growing a thin layer of Cesium that lowers the bulk conduction band.

The extracted electrons have the same helicity as the incident photons since they exit in the direction opposite to that of the photons. The maximum polarization of 100% could be achieved from a strained GaAs cathode, compared to regular GaAs which is limited to 50% polarization due to degeneracy of $J = 3/2$ state. In practice, polarizations of up to 85% were achieved from the strained cathode.

After photoemission the electrons were accelerated across the 60 kV potential of the gun, creating a 200−250 ns pulse of $\approx 0.5-2\times10^{11}$ electrons which were bunched and accelerated down the linac. The source operated at 120 Hz, but typically only 119 were used by E154. One in every 120 pulses was a short, high-intensity 3 ns pulse used for accelerator diagnostics and tuning. This *"witness" pulse* was deflected in the Beam Switchyard and did not enter the End Station A.

2.2.2 Beam acceleration and transport

The electrons created at the source entered the 3.2 km linear accelerator. The accelerator structure consists of 30 sectors, each containing eight klystrons, steering dipoles, a quadrupole, and additional elements for monitoring the beam position and current.

Each of the 240 klystrons feed microwave radiation at 2856 MHz into copper cavities. During E154 (and during the SLC running) a technique called SLAC Energy Doubling (SLED) was used that involves reversal of the microwave phase during the pulse. This results in a sharp increase in the accelerating voltage in the cavities at the expense of reducing the pulse length to $200 - 300$ ns (compared to ≈ 2 μs in the normal, non-SLED mode). The maximum beam energy achievable in the SLED mode is ≈ 52 GeV.

At the end of the linac, the beam is deflected by $0.5°$ into the A-line by two magnets in the Beam Switchyard (BSY). The A-line consists of twelve identical 3 m long dipole magnets operated in series, each of which bends the beam by $\approx 2°$ from the linac towards End Station A (ESA). Twelve quadrupoles are used to control the beam divergence and the spot size at the target.

2.2.3 Spin precession and beam energy

The helicity of electrons entering the linac is the same as that of photons hitting the photocathode and it remains constant during acceleration to the end of the linac. Two dipole magnets located in the BSY deflect the beam by $0.5°$ north into the A-Line where the set of twelve dipole magnets bend it by 24.0° north into the End Station A. During deflection, the spin of the electrons precesses by an amount

$$\Delta\theta_{\text{precession}} = \gamma\Delta\theta_{\text{bend}}\left(\frac{g-2}{2}\right) \tag{2.1}$$

where $\gamma = E/m$, $\Delta\theta_{\text{bend}} = 24.5°$, and g is the gyromagnetic ratio of the electrons.[20] For beam energies which are even multiples of 3.2374 GeV, the electron helicity at the target in ESA is the same as it is at the source. E154 ran at the linac energy of 48.56 GeV which is an odd multiple, and the helicity of the electrons at the target was opposite to that at the source.

At high beam energies, synchrotron radiation becomes important. In general, the energy loss due to the synchrotron radiation is given by

$$\Delta E_{\text{synch}} = \frac{2\theta_{\text{bend}}}{3}\frac{e^2}{\rho}\left(\frac{E}{m}\right)^4 \tag{2.2}$$

where $\rho = 85.927$ m is the bending radius. Taking into account the synchrotron energy loss, the "magic energy" corresponding to 15π spin rotations is $E = 48.362$ GeV. Experiment E154 took polarized data at a beam energy $E = 48.325$ GeV with an uncertainty on the value of ≈ 0.05 GeV.[90] It was determined by measuring the energy dependence of the beam polarization in the Møller polarimeter.[91]

Synchrotron radiation also results in the growth of the beam emittance and therefore a larger spot size at the target. In order to reduce the spot size to the required ≈ 1 mm, a quadrupole (Q41) was placed in the alcove of the ESA. The focal point was at the target, and the beam gradually diverged after the target.

2.2.4 Electron beam helicity reversal

The ability to reverse the beam polarization on a pulse-to-pulse basis with the Pockels cell was very important for reducing systematic errors. Possible false asymmetries due to slow changes in spectrometer acceptance and detector efficiencies were averaged out by the rapid beam helicity reversals. Also, by changing the target polarization direction periodically, we could further reduce the effect of asymmetries induced by possible helicity-dependent differences in the beam properties. The absolute helicity of the light was determined by the sign of the asymmetry measured by the Møller polarimeter.

The beam helicity was chosen on a pulse-to-pulse basis using a pseudo-random bit generator. The sequence began when a 32 bit random number was chosen, with the last (32nd) bit determining the helicity of the beam. For the next pulse, the 19th and 32nd bit determined the new first bit via a logical XOR, and the remaining bits were shifted up. The new 32nd bit determined the helicity of this pulse, and the cycle was repeated. The sequence of the 33 consecutive polarization bits thus determined the polarization state of all consecutive spills. The generator was running continuously even when the source was off. Thus, we could predict the polarization state of each spill knowing the spill number and could test the integrity of the system.[92]

The polarization state of the beam was sent to the Data Acquisition System via four physically distinct pathways: the PMON line, Mach line, Pockels Cell High Voltage Line, and the Veto Bits. Each pathway delivered a two-bit combination, where the combination '01' referred to positive helicity photons and '10' meant that the photons incident on the photocathode were of negative helicity. Combination '00' in the PMON meant that the beam is unpolarized, and '11' marked an error condition. Typically, all five (four hardware signals and the predictor) methods of determining the polarization state of the spill agreed[92] and we discarded the pulses if a disagreement was detected. If the rate of mismatches exceeded a conservative limit of $5 \cdot 10^{-4}$, the entire run was discarded (see Section 3.7.2).

2.3 Beam monitoring

2.3.1 Beam energy

The energy of the beam was monitored continuously and recorded at every checkpoint (typically, three or four times per half an hour run). A flip-coil was located in an off-line dipole, identical to the twelve used in the A-line and placed in series with them. The current induced in this flip-coil is proportional to the magnetic field in the dipoles. In turn, the magnetic field strength determines the momentum of the electrons passing through the A-line to about 0.1%. The absolute energy scale was also determined independently by measuring the energy dependence of the beam polarization (see below). The energy spread of the beam, $\Delta E/E$, entering ESA was restricted to 1% full width by using three adjustable slits located in the A-line.

2.3.2 Beam current

On a pulse-to-pulse basis the beam charge was measured with two independent ferromagnetic toroids. One was located about 38 m upstream of the target, and the second a few meters upstream. As the beam passed through the ferrite core of the toroid, a signal was induced in loops of wire wrapped around the toroid, which in turn was sent to a resonant LC circuit. The induced current was amplified and integrated, and hence the total charge was proportional to the beam current. Each toroid was calibrated by discharging a precisely charged capacitor through the toroid and measuring the induced charge. The calibration coefficients were extracted several times a day for each toroid, and the total beam charge of each spill was measured to about 0.5%.

2.3.3 Beam position and steering

The beam position and width were monitored with a wire array positioned 10.5 m downstream of the target. Each of the 24 wires in both (x and y) projections were made of 0.127 mm diameter CuBe, with a spacing of 1.1 mm. The wire arrays were read out on a pulse-to-pulse basis by the beam branch of our Data Acquisition System (DAQ); this information was also available to linac operators at the Main Control Center (MCC) to perform minor steering corrections to the beam. Two mylar roller screens, coated with fluorescent ZnS and observed with television monitors, were placed periodically into the beam and allowed a coarse centering and focusing of the beam. The first screen was placed ≈ 25 m upstream from the target right after the quadrupole Q41. The second roller screen was in front of the target.

Additional information about the beam quality came from two scintillator counters. The first one (dubbed a *"bad spill monitor"*) was located upstream of the target in the End Station alcove and was sensitive to beam scraping. The second counter (a *"good spill monitor"*) was downstream of the target, several meters away from the beam pipe. It was sensitive primarily to particles created at the target and served as a good indicator of the time structure of the beam. Signals from both counters were displayed on oscilloscopes (both in the Counting House and MCC) and monitored continuously. The signals were also integrated by ADCs and stored on tape to be used in the data analysis.

Other monitors of the beam included two traveling wave beam position monitors (TWBPM) located in front of the target. One cavity produced an RF signal proportional to the deviation of the beam from the horizontal center of the cavity, the second was sensitive to vertical displacement. A final system used for the target protection was a secondary emission monitor (SEM) made of a thin aluminum foil with a circular aperture of 2 cm for the beam that matched the diameter of a target cell. It was placed immediately upstream of the target. Large signals from the SEM indicated a potentially destructive displacement of the beam, prompting an automatic beam shutdown. This prevented the beam from hitting the thick side walls of the target.

2.4 Møller polarimeter

The beam polarization was measured periodically by a single arm Møller polarimeter. The polarimeter was located upstream of the target, and consisted of the polarized iron foil target, momentum-analyzing magnet, and a set of finely-segmented silicon detectors. The polarimeter was also used to optimize the beam energy and the source laser wavelength with respect to the beam polarization.

2.4.1 Møller asymmetry

The cross section for polarized e^-e^- scattering is fully calculable in QED, and in the center-of-mass frame is given by[93]:

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{s}\frac{(3+\cos^2\theta)^2}{\sin^4\theta}[1 + P_z^B P_z^T A_{zz}(\theta)] \tag{2.3}$$

for longitudinally-polarized beam and target, where the asymmetry is

$$A_{zz}(\theta) = -\frac{(7+\cos^2\theta)\sin^2\theta}{(3+\cos^2\theta)^2} \,. \tag{2.4}$$

Here s and θ are the CM energy squared and the scattering angle, P_z^B and P_z^T are the beam and target foil longitudinal polarizations, respectively. By measuring the Møller scattering rate for the electron beam and target spins aligned, $\sigma^{\uparrow\uparrow}$, and anti-aligned, $\sigma^{\uparrow\downarrow}$, we can form the asymmetry

$$A = \frac{\sigma^{\uparrow\uparrow} - \sigma^{\uparrow\downarrow}}{\sigma^{\uparrow\uparrow} + \sigma^{\uparrow\downarrow}} = P_z^B P_z^T A_{zz}(\theta). \tag{2.5}$$

If the target polarization is independently known, a measurement of the Møller asymmetry determines the beam polarization.

The asymmetry maximum is at $\theta_{CM} = 90°$ where the unpolarized laboratory cross section is 0.179 b/sr and $A_{zz} = -7/9$. With a typical foil polarization of 0.08 (that corresponds to 2 outer-shell electrons of Fe aligned with the external magnetic field) and beam polarization of 0.8, the expected asymmetry is roughly 0.05.[91]

2.4.2 Polarimeter apparatus

The polarimeter consisted of polarized target foils, a mask to define the azimuthal and vertical acceptance, a magnet to analyze the momentum of the scattered electrons, and detectors to measure the scattering rate. The mask selected Møller scattered electrons in the vertical plane and the magnet deflected them horizontally. The beam vacuum pipe went through the magnet and was enclosed in an iron septum to eliminate the field inside the beam pipe. The elastically-scattered Møller electrons have correlated momenta and scattering angles, and form a stripe at the plane of the detector. The top silicon detector contained 48 channels placed on two 4×6 cm^2 pads to produce a segmentation of 2.18 mm in the vertical (θ) direction. It was mounted on a remotely-controlled mover and could be positioned anywhere within the Møller acceptance. Four separate silicon detectors with coarser pitch were mounted side by side at fixed positions in the bottom aperture.

The E154 polarimeter was an evolution of the previous End Station A Møller Polarimeters[94] and utilizes many ideas and components of previous designs. The top and side views of the polarimeter are shown in Fig. 2.3.

2.4.3 Foil polarization

The foil polarization was determined by measuring its magnetization $\vec{\mathbf{M}}$.[95] Ramping the external field $\vec{\mathbf{H}}$, induces a voltage in a pickup coil wound around the foil $\int V dt = \Phi_f - \Phi_i$. The flux Φ has contributions from the foil and air:

$$\Phi = N_{\text{turns}} \left[\vec{\mathbf{B}}_{\text{foil}} \cdot \vec{\mathbf{A}}_{\text{foil}} + \vec{\mathbf{H}} \cdot (\vec{\mathbf{A}}_{\text{coil}} - \vec{\mathbf{A}}_{\text{foil}}) \right] \ , \tag{2.6}$$

where $N_{\text{turns}} = 500$ is the number of turns of the pickup coil. By integrating the induced voltage with and without the foil, and noting that $\vec{\mathbf{B}}_{\text{foil}} = \vec{\mathbf{H}} + 4\pi\vec{\mathbf{M}}$, we obtain

$$4\pi|\vec{\mathbf{M}}| = \frac{1}{N_{\text{turn}}|\vec{\mathbf{A}}_{\text{foil}}|} \left[\int_{\text{foil in}} V dt - \int_{\text{foil out}} V dt \right] \tag{2.7}$$

Fig. 2.3. Top (top picture) and side (bottom picture) views of the E154 Møller Polarimeter.

The magnetization is related to the foil polarization by

$$P_z^T = \left[\frac{g}{g-1}\frac{g'-1}{g'}\right]\frac{|\vec{\mathbf{M}}|}{n\mu_B} \tag{2.8}$$

where g is the gyromagnetic ratio of the electron,[20] $g' = 1.916 \pm 0.02$ is the gyromagnetic ratio of Permendur,[96] n is the number density of electrons, and μ_B is the Bohr magneton.

Six foils of approximate thicknesses 20, 20, 30, 40, 40, and 154 μm were installed at a 20.7° relative to the beam line. The majority of E154 Møller data were taken with a 40 μm foil. The foils were magnetized to saturation by $\approx$ 100 G field produced by Helmholtz coils. The polarity of the coils was typically reversed between the Møller runs to reduce systematic effects. The magnetization of the foils was measured before and after E154. Both measurements agreed within 0.6% with an average change of less than 0.1%. A relative systematic error of 1.9% was assigned to the foil polarization.

2.4.4 Beam polarization results of the Møller runs

Møller data were taken periodically throughout the experiment for a total of about 140 runs. The signals from the silicon detectors were pre-amplified and read out by the ADCs. In each run, the average pulse heights in every channel were recorded for the two beam helicities, from which the (R–L) and (R+L) line-shapes were formed. The background under the unpolarized (R+L) Møller line-shape was estimated by fitting the (R+L) line-shape to a quadratic polynomial plus the line-shape expected from unpolarized Møller scattering.[91] The unpolarized Møller line-shape was obtained by using the (R–L) shape with the corrections for the atomic motion of target electrons,[97] and was in a very good agreement with a Monte Carlo calculation. The (R–L) and (R+L) line-shapes were integrated and the Møller asymmetry was formed. The background subtraction increased the measured asymmetry by 17–24%. The statistical errors on the beam polarization were typically

Table 2.1. Measured longitudinal beam polarization for different running periods. The error is statistical only.

Run range	Polarization
1329–1411	0.759 ± 0.004
1456–1684	0.775 ± 0.005
1691–2311	0.814 ± 0.002
2316–3371	0.824 ± 0.001
3377–3788	0.826 ± 0.002

Table 2.2. Systematic error contributions to the beam polarization measurement.

Systematic error contribution	Value
Foil magnetization	1.9%
Kinematic acceptance	0.3%
Background correction	2.0%
Fit range	0.3%
TOTAL	2.8%

$0.003 - 0.006$ per run. The history of the beam polarization measured by the polarimeter is shown in Fig. 2.4 and is given in Table 2.1. The average beam polarization (weighted by statistics of $A_{\|}$) was 81.8%.

2.4.5 Systematic errors

The overall systematic error has contributions from the foil polarization, uncertainties in the expected Møller asymmetry for each detector, and uncertainties in the background subtraction. The various contributions to the systematic error are summarized in Table 2.2.[91]

2.5 Polarized ^{3}He target

The polarized ^{3}He target was one of the major factors that determined the success of this experiment. It was very similar to the polarized target used in E142.[98] The target was a two-chambered 30 cm long glass cell containing ^{3}He at densities of



Fig. 2.4. Measured longitudinal beam polarization versus E154 run number. Errors are statistical only. "Jumps" in the measured value of polarization indicate various adjustments made to the source during the run.

2.6×10^{20} atoms/cm^3. The ^{3}He nuclei were polarized in the spin exchange collisions with optically pumped Rubidium atoms. Target polarizations of nearly 50% were achieved with an average polarization of 38%.

2.5.1 Optical pumping and spin exchange

Optical pumping is an effective technique for achieving high polarizations in a high density noble gas target.[99] The process is illustrated in Fig. 2.5. For example,for a positive helicity, circularly polarized laser light at 794.7 nm excites the $5S_{1/2}(M_J = -1/2)$ ground state to the $5P_{1/2}(M_J = +1/2)$ in Rubidium vapor (the D1 line). Radiative decays to the ground state favor the $5S_{1/2}(M_J = -1/2)$ state over $5S_{1/2}(M_J = +1/2)$ state by a factor two. By adding 65 torr of N_2, as



Fig. 2.5. Schematic of the optical pumping and spin exchange processes.

in our target cells, the $5P_{1/2}(M_J = \pm 1/2)$ states are collisionally mixed at a rate exceeding the radiative decay. The relaxation into the $5S_{1/2}$ ground states occurs equally into the two sub-states, and only two photons are required to increase the $5S_{1/2}(M_J = +1/2)$ population by one. The polarization of the valence electrons of Rb competes with several depolarizing mechanisms, the most of important of which are from Rb-Rb and Rb-^{3}He collisions (the latter is dominant at high pressures), as well as Rb-target cell wall interactions. However, with sufficiently high Rb vapor density (10^{14} atoms/cm^3) and high laser power, the optical pumping rate of $1/\gamma_{\rm pump} \approx 10^{-6}$ s exceeds the $1/\gamma_{\rm relax} \approx 10^{-3}$ s relaxation rate, and almost 100% Rubidium polarization could be achieved.

The polarization of the Rb atoms is transferred to the ^{3}He nuclei in the spin exchange collisions due to the hyperfine interaction of the Rb electron magnetic moment and the magnetic moment of the ^{3}He nucleus, as shown in Fig. 2.5. The spin exchange rate, γ_{SE}, is proportional to the relative velocity between the ^{3}He and Rb, v, and to the number density of Rb, [Rb]:

$$\gamma_{SE} \equiv <\sigma_{SE} v> [\mathrm{Rb}] \tag{2.9}$$

where the velocity averaged cross section is $\approx 1.2 \times 10^{-19}$ cm^2/s.[100] With [Rb] $\approx 10^{14}$ cm^{-3}, polarization times for ^{3}He are very long, $1/\gamma_{SE} \approx 25 - 40$ hours.

Various processes are responsible for depolarization of the ^{3}He atoms. Collisions between ^{3}He atoms can cause the exchange of the nuclear spin and orbital angular momentum. Paramagnetic impurities, both gaseous and embedded in the cell walls, contribute further to the spin relaxation. Factors external to the cell also play a role. Polarized ^{3}He diffusing through regions in which the holding field is inhomogeneous, precesses about the local field and the spin direction is randomized. This effect is usually minimized by keeping the gradients perpendicular to the alignment field small ($<$ 20 mG/cm). Nuclear relaxation is also induced by the ion production in the cell by the electron beam due to creation of non-zero electron spin states in the ionized ^{3}He atom.[101] The cumulative effect of these relaxation processes can be characterized by

$$\Gamma_R = \frac{1}{\tau_{\mathrm{cell}}} + \frac{1}{\tau_{\nabla B}} + \frac{1}{\tau_{\mathrm{beam}}}\,, \tag{2.10}$$

where τ_{cell}, $\tau_{\nabla B}$, and τ_{beam} are intrinsic cell relaxation time, and relaxation times induced by the magnetic field gradients and the electron beam. Typical relaxation times for our target cells were on the order of $40 - 60$ hours.

The time dependence of the ^{3}He polarization is given by a simple model of the competing spin exchange and relaxation processes

$$P_{^3\mathrm{He}}(t) = \frac{\gamma_{SE}}{\gamma_{SE} + \Gamma_R} P_{\mathrm{Rb}} \left[1 - \exp\left(-(\gamma_{SE} + \Gamma_R)t\right)\right]\,, \tag{2.11}$$

assuming $P_{^3\mathrm{He}}(t=0) = 0$, and where P_{Rb} is the equilibrium polarization of Rb $\approx$ 100%. The polarization buildup ("spin-up") observed in one of the best cells, Picard, is shown in Fig. 2.6. The maximum polarization of 47.9% was achieved.



Fig. 2.6. The on-line plot of the polarization buildup in the target cell Picard. The calibration constant was reduced by $\approx 2\%$ in the post-run analysis.

2.5.2 Target apparatus

The target cell was made of Corning 1720 aluminosilicate glass, which has low ^{3}He permeability. It used a two chamber design where the upper, pumping cell with a volume of $70-80$ cm^3 contained ^{3}He and a few tens of mg of Rb. This chamber was enclosed in a plastic oven and heated to 170-200° C to create a Rb vapor density of about 10^{14} atoms/cm^3. Three 20 W diode lasers and four Argon-ion pumped Ti:sapphire lasers were located in a specially constructed laser hut near the target. The lasers pumped the target continuously, resulting in a total of almost 80 watts of power[1] on the pumping cell. The light from each laser passed through the various optical components including a quarter-wave plate to produce circularly polarized light, before reaching the pumping cell which was about 3 m away. A schematic of the target is shown in Fig. 2.7.

[1]The spectrum of the diode lasers is quite broad so the total power absorbed by Rb at 795 nm was somewhat smaller.



Fig. 2.7. The schematic of the E154 polarized target setup.

The ^{3}He atoms polarized in the upper cell diffused through a 60 mm long transfer tube to the lower, target cell, with a time constant of about 10 minutes. This cell was a cylinder $\approx$ 30 cm long, 1 cm in radius, with $\approx$ 50 μm thick endcaps of concave shape. Since the glass is stronger under compression than under tension, the concave design allowed thinner windows than the traditional convex, shape for the same ^{3}He pressure. E154 benefited greatly from this feature since the dilution factor (the scattering rate from polarized ^{3}He relative to the total rate) was increased by almost a factor of 2 compared to E142. Cooling jets of ^{4}He were directed at the endcaps to relieve the heating caused by the beam. The target cell temperatures were monitored with 6 Resistive Thermal Devices (RTDs) mounted in various places on the target and pumping cells. The temperature throughout the target cell remained at $60-80$ °C (depending on the position). The residual Rb density was of the order 10^{11} atoms/cm^3, which is insignificant in comparison to the 2.6×10^{20} atoms/cm^3 of ^{3}He and the 2.4×10^{18} molecules/cm^3 of N_2 which were the primary components of the target cell. The target holder had three positions with the reference cell placed below the polarized cell (the third position was empty). The reference cell was used for the experimental studies of the dilution factor (see Section 3.10), and the "no target" runs were used to monitor the beam halo[2]. The precision mover placed either of two cells into the beam with a sub-millimeter accuracy. A large scattering chamber enclosed the target assembly, and was kept under vacuum of a few mtorr.

Outside of the scattering chamber was a set of 1.4 m diameter Helmholtz coils which produced a 10–30 G holding field to align the nuclear spins of ^{3}He. A similar set of coils perpendicular to the beam direction was used to achieve the transverse polarization for measurements of $A_\perp$.

[2]The initial estimates of the beam halo were obtained with a special "dummy" cell that had no windows. Since such a cell was not in the target holder during most of the run, results of the "no target" runs were used for monitoring the halo, and the "dummy" cell runs were used as a calibration.

2.5.3 Target cells

The thin target windows could not sustain the high intensity electron beam for more than a week. Several effects could be blamed for breaking of the windows, but the exact cause is unknown. A total of nine polarized target cells and four reference cells were used in the experiment. The catalog of the cells with the relevant dimensions is given in Table 2.3.[102]

The target windows thickness was measured by three independent methods: by a micrometer, by X-ray absorption (using ^{55}Fe source), and using the interference of the laser light reflected from two surfaces of a window. The optical method is the most precise and gives an uncertainty of 3%, limited by the non-uniformity of the windows. The windows that broke in the beam were not measured by the optical technique, and the uncertainty in their thicknesses is 5%.

The ^{3}He density was measured during the filling of the cell, and later using the pressure broadening of the Rb absorption line. The fill density was corrected for the difference in the pumping and target cell temperatures. The error on the density is 2%.

2.5.4 Target polarimetry and polarization results

The ^{3}He polarization was determined by two independent techniques.[103] The traditional method, the adiabatic fast passage (AFP) NMR, was used during the run.[104] A set of 42.75 cm diameter Helmholtz drive coils above and below the target provided a 72 mG RF field at 92 kHz, while the main holding field was swept from 18 G to 36 G, through the Larmor resonance at 28.4 G. The field sweep rate was optimized to minimize the polarization losses. The rate of 1.2 G/s was slow enough for the nuclear spins to follow the changing field, but fast enough to avoid the de-phasing of the spins while passing through the resonance. The resulting nuclear spin flip induced a signal in a set of 200 turn copper pickup coils, wound

Table 2.3. Target cells used in E154.

Cell Name	Runs	Windows (μm)	Length (mm)	^{3}He density (10^{20} cm^{-3})	N_2 density (10^{18} cm^{-3})	Fate
Dave	1201 – 1388	45.0, 53.5	291	2.70	2.3	Broke
Riker	1412 – 1757	66.5, 60.0	300	2.68	1.4	Broke
Bob	1777 – 2043	70.1, 69.7	299	2.68	2.3	Broke
SMC	2050 – 2311	68.8, 64.2	302	2.68	2.4	Survived
Generals	2316 – 2594	52.2, 56.7	297	2.63	2.4	Broke
Hermes	2597 – 2902	46.9, 56.6	295	2.64	2.4	Broke
Prelims	2903 – 3100	45.1, 42.2	300	2.68	2.3	Broke
Chance	3101 – 3371	83.0, 83.0	299	2.68	2.3	Survived
Picard	3377 – 3788	69.3, 61.6	299	2.64	2.3	Survived
Ref. Cell #1	733 – 1757	58.4, 46.5	304			Broke
Ref. Cell #2	1777 – 2047	73.5, 46.3	297			Survived
Ref. Cell #3	2050 – 2596	50.9, 48.5	304			Survived
Ref. Cell #4	2597 – 3788	70.8, 64.0	302			Survived



Fig. 2.8. Typical ^{3}He and water AFP-NMR signals.

on a 76.2 mm by 25.4 mm Teflon form and centered around the target cell. Using a 100 pF capacitor, an LC resonant circuit was formed. The signal was pre-amplified, then went through a lock-in amplifier that subtracted the 92 kHz modulation. The resulting NMR signal was proportional to the ^{3}He polarization. The proportionality constant was determined from the thermal equilibrium Boltzmann polarization of protons in a water sample. The water signals were 10^5 times smaller compared to the ^{3}He signals, so typically many sweeps were averaged. Typical ^{3}He and water signals are shown in Fig. 2.8.

The primary sources of the systematic uncertainty in the AFP-NMR polarimetry were the height of the water signal (1.8%), proton polarization in water (1.0%), and ^{3}He density (1.9%). The total systematic error of the AFP method is 3.4%.[103]

The second method, used to calibrate the AFP-NMR polarization measurement, was based on the Electron Paramagnetic Resonance (EPR) technique.[105] It used



Fig. 2.9. EPR frequency shift due to reversal of ^{3}He polarization.

the shift of the Rb Zeeman resonance due to the magnetic field created by the polarized ^{3}He. The shift is quite substantial (about 20 kHz out of 8 MHz) and easy to measure. If the Rb polarization is very high, most of the atoms are in $F = 3$, $M = \pm 3$ state (where F and M are the total angular momentum and its projection for the nuclear and electron spins of ^{95}Rb). By applying the RF frequency, one can induce the transitions between neighboring M sublevels of the $F = 3$ manifold. The EPR resonance is observed by monitoring the rate of fluorescence at the D_2 Rubidium line as a function of RF frequency.[103] The RF field was created by a coil mounted on a side of the oven. The fluorescence was detected by a photodiode with a D_2-line filter. The magnetic field was measured to one part in 10^5 by a Flux-Gate magnetometer. The frequency shift due to ^{3}He polarization was detected by reversing the ^{3}He polarization. A typical EPR signal is shown in Fig. 2.9. The quality of the data is very good and the size of the frequency shift was extracted with an error of less than 0.5%.

The frequency shift is proportional to the ^{3}He polarization where the coefficient of the proportionality κ_0 is specific to the Rb-^{3}He system.[105] This coefficient was measured[103] with an uncertainty of 1.0%. The errors in the EPR measurement mostly came from κ_0, the polarization gradient between the pumping (where the polarization is measured by EPR) and the target cells (1.5%), and the spread of the data (1.3%). The total systematic error of the measurement is 3.0%.

The AFP and EPR polarization measurements disagreed by 5.4%. If the uncertainties of each measurement are treated as independent and the shapes of probability distributions are Gaussian, the probability of such a disagreement is 23%. However, an unknown systematic effect could alter one of the measurements. We therefore increase the polarimetry error so that it covers both measurements and their errors. The final error also includes a 1.7% drift in AFP calibration constants measured before and after the run. The final target polarimetry error is 4.8%.

2.5.5 ^{3}He polarization direction

The direction of the ^{3}He polarization relative to the magnetic field was determined in three ways.[106] The first method used the sign of the AFP-NMR signals. Protons and ^{3}He have magnetic moments of opposite sign, so if their spins are in the same directions initially and are swept through a resonance with $d|\mathbf{B}|/dt$ of the same sign, the NMR resonance signals are of opposite sign. This determines the ^{3}He polarization direction with respect to the proton, which is preferentially aligned parallel to the external fields. The second method relates the sign of the EPR frequency shift which is proportional to the ^{3}He polarization. The third method was based on the observation of the masing effect during AFP due to the coupling of ^{3}He spins to the pickup coils. All three methods were consistent.

The direction of the holding field was determined by a flux-gate magnetometer, by a Hall probe relative to the spectrometer magnets, by compass, and finally by the direction of the current in the Helmholtz coils. Again, all methods were consistent. In addition, liquid crystal circular polarizers determined the laser helicity, which determines the angular momentum transferred to the ^{3}He nuclei, and hence the polarization direction.

2.6 Magnetic spectrometers

2.6.1 Introduction

Design of the magnetic spectrometers for the 50 GeV spin structure program at End Station A was driven by the physics requirements. The goal of the experiments is to measure the spin structure functions of the neutron and proton down to the lowest possible x and over a more extensive Q^2 range than compared to the previous ESA experiments.[9–11] Design choices reflect the requirements of high counting rates, while keeping the ability to unambiguously identify scattered electrons and determine their energy in the presence of large background of charged hadrons and low energy neutral particles. An important factor is a short SLED beam pulse, which leads to a factor of $10 - 20$ increase in the instantaneous rates compared to E142 and E143 (if the rate per spill is kept the same).

The system consists of two independent magnetic spectrometers centered around $2.75°$ and $5.5°$ relative to the beam direction. The 48.3 GeV beam combined with the choice of angles allows the measurements in the kinematic range $0.014 \leq x \leq 0.8$ and $1 \text{ GeV}^2 \leq Q^2 \leq 17 \text{ GeV}^2$ (Fig. 2.10). This is the most extensive kinematic range ever achieved by polarized electron scattering experiments to date. The $2.75°$ spectrometer covers a momentum range from 10 to 44 GeV, and the $5.5°$ spectrometer covers a momentum range from 10 to 39 GeV. The low momentum cutoff is dictated by the requirement of $y = (E - E')/E < 0.8$ imposed by radiative corrections (which increase rapidly at high y), and the desire to keep the charged hadron background at a tolerable level. The high momentum cutoff is set by the standard deep-inelastic cut $W^2 > 4 \text{ GeV}^2$. The theoretically calculated differential cross sections $d^2\sigma/d\Omega dE$ for the two spectrometers are plotted in Fig. 2.11.

A system of two independent "closed-geometry" magnetic spectrometers has several advantages. First, the neutral background is highly suppressed with the so-called *"double-bounce"* geometry that prevents neutral particles from reaching the