

Introduction



What is parity violation?

The parity symmetry states that the physical laws behind all natural phenomena must be the same as those behind their mirror images.


It has been long confirmed that parity is conserved in strong and electromagnetic interactions. However, the postulate that parity is violated in weak interactions was proposed by Lee and Yang in 1956, and was soon confirmed experimentally by Wu.



Tsung-Dao Lee




Chen-Ning Yang



Chien-Shung Wu

A polarized electron and its mirror image




What is PV Deep Inelastic electron Scattering (PVDIS)?

For polarized electrons, the mirror image of a left-handed electron is right-handed;

There is a difference in scattering probability between left- and right-handed electrons, causing a cross-section asymmetry. This asymmetry comes from the interference term between the photon- and the Z^0 -exchanges.

In deep inelastic scattering, the internal structure of the target nuclei or nucleon is revealed at the quark and gluon level.



PVDIS asymmetry from a deuterium target:

$$A_d = (540 \text{ ppm}) Q^2 \frac{2C_{1u}[1+R_c(x)] - C_{1d}[1+R_s(x)] + Y(2C_{2u} - C_{2d})R_v(x)}{5+R_s(x)+4R_c(x)}$$

Q^2 : Four-momentum transfer squared between the electron and the target nucleon

x : variable relevant to the nucleon internal structure

$$C_{1u} = g_A^e g_V^u = -\frac{1}{2} + \frac{4}{3} \sin^2(\theta_w)$$

$$C_{2u} = g_V^e g_A^u = -\frac{1}{2} + 2 \sin^2(\theta_w)$$

$$C_{1d} = g_A^e g_V^d = \frac{1}{2} - \frac{2}{3} \sin^2(\theta_w)$$

$$C_{2d} = g_V^e g_A^d = \frac{1}{2} - 2 \sin^2(\theta_w)$$

In the SM to the first order

From A_d one can extract $C_{1,2q}$ and $\sin^2\theta_w$, providing a test of the electroweak (EW) standard model (SM).

$R_c(x)$, $R_s(x)$, $R_v(x)$ are related to parton distribution functions, thus A_d is sensitive to internal structure of the nucleon, which is described by QCD – the SM of strong interactions.

In 1972, PVDIS result from SLAC E122 was consistent with $\sin^2\theta_w=1/4$, confirmed the Standard Model prediction; However, PVDIS measurement has not been performed again.

Precision Measurement of Parity-Violating Asymmetry in Deep Inelastic e^2H Scattering (PVDIS) Using a 6 GeV Beam at JLab

— Low energy tests of the Standard Model of particle physics in the electroweak and hadronic sectors

Xiaochao Zheng (UVa), Robert Michaels (JLab), Paul Reimer (ANL), Ramesh Subedi (UVa)



JLab Experiment 05-007

The ultimate, long term goal of the JLab PVDIS program is to study both the Standard Model of EW interactions and hadronic phenomena.

Our first step is to use a 6 GeV beam -- JLab E05-007 (planned for 2009):

- Use 85 μ A, 6 GeV, 80% polarized beam on a 25-cm LD2 target;
- Two Hall A High Resolution Spectrometers detect scattered electrons;
- Measure A_d at $Q^2=1.10$ and 1.90 GeV^2 to about 2% (stat.), with similar systematic uncertainties.


Assuming hadronic effects are small and using the expected known values for C_{1q} 's, we can extract $2C_{2u}-C_{2d}$; Will also provide crucial guidance to the 12 GeV program.



Experimental upgrade needed:


- upgrade from IR to green laser to provide 1% precision

- upgrade to flash-ADC based DAQ to count at $\sim 1\text{MHz}$ rates



Electronics being assembled at JLab


The new electronics will be installed in parallel to the regular DAQ



PVDIS at JLab 12 GeV


Provide precision test of the Electroweak Standard Model, complementary to other experiments:

- High-energy direct searches (such as LHC) may discover new particles. However to study their coupling properties we must use low- and medium energy experiments;
- Moller (SLAC E158, completed): pure-leptonic;
- Qweak (JLab Hall C, future): only C_{1q} 's;
- Only PVDIS is sensitive to C_{2q} 's (new physics in the quark axial coupling).




Expected results on $2C_{2u}-C_{2d}$ from PR12-07-102, assuming small hadronic effects

Expected results on $\sin^2\theta_w$ from PR12-07-102, assuming small hadronic effects



Expected constraint on CSV from PR12-07-102




Provide possible direct observation of:

- Non-perturbative QCD effects in the EW sector;
- Charge symmetry violation at the quark level.

(All these are not yet calculable in QCD ! Direct observations will be very exciting.)

References

- R. Michaels, P.E. Reimer, X. Zheng, et al, JLab Proposal PR05-007.
- K. Paschke, P.E. Reimer, X. Zheng et al., JLab 12 GeV Proposal PR12-07-102.
- JLab 12 GeV electroweak physics meeting, Initiative on the large acceptance solenoid device, October 11, 2007.



Jefferson Lab Experimental Hall A