Dynamical Relativistic Effects Observed in Quasielastic 1p–Shell Proton Knockout from ^{16}O

J. Gao,[17] B. D. Anderson,[13] K. A. Aniol,[2] L. Auerbach,[29] F. T. Baker,[7] E. J. Beise,[16] J. Berthot,[1] W. Bertozzi,[17] P. -Y. Bertin,[1] L. Bimbot,[22] W. U. Boeglin,[5] E. J. Brash,[24] V. Breton,[1] H. Breuer,[16] E. Burtin,[26] J. R. Calarco,[18] L. Cardman,[30] G. D. Cates,[23] C. Cavata,[26] C. C. Chang,[16] N. S. Chant,[16] J. -P. Chen,[30] E. Cisbani,[12] D. S. Dale,[14] R. De Leo,[10] A. Deur,[1] B. Diederich,[21] P. Djawotho,[33] J. Domingo,[30] B. Doyle,[14] J. -E. Ducret,[26] M. B. Epstein,[2] L. A. Ewell,[16] J. M. Finn,[33] K. G. Fissum,[17] H. Fonvieille,[1] B. Frois,[26] S. Frullani,[12] F. Garibaldi,[12] A. Gasparian,[8,14] S. Gilad,[17] R. Gilman,[25,30] A. Glamazdin,[15] C. Glashausser,[25] J. Gomez,[30] V. Gorbenko,[15] T. Gorringe,[14] K. Griffioen,[33] F. W. Hersman,[18] R. Holmes,[28] M. Holtrop,[18] N. d'Hose,[26] C. Howell,[4] G. M. Huber,[24] C. E. Hyde-Wright,[21] M. Iodice,[12] C. W. de Jager,[30] S. Jaminion,[1] M. K. Jones,[33] K. Joo,[32] C. Jutier,[1,21] W. Kahl,[28] S. Kato,[34] J. J. Kelly,[16] S. Kerhoas,[26] M. Khandaker,[19] M. Khayat,[13] K. Kino,[31] W. Korsch,[14] L. Kramer,[5] K. S. Kumar,[23] G. Kumbartzki,[25] G. Laveissière,[1] A. Leone,[11] J. J. LeRose,[30] L. Levchuk,[15] M. Liang,[30] R. A. Lindgren,[32] N. Liyanage,[17] G. J. Lolos,[24] R. W. Lourie,[27] R. Madey,[8,13,30] K. Maeda,[31] S. Malov,[25] D. M. Manley,[13] D. J. Margaziotis[2] P. Markowitz,[5] J. Marroncle,[26] J. Martino,[26] J. S. McCarthy,[32] K. McCormick,[21] J. McIntyre,[25] R. L. J. van der Meer,[24] Z. -E. Meziani,[29] R. Michaels,[30] J. Mougey,[3] S. Nanda,[30] D. Neyret,[26] E. A. J. M. Offermann,[30] Z. Papandreou,[24] C. F. Perdrisat,[33] R. Perrino,[11] G. G. Petratos,[13] S. Platchkov,[26] R. Pomatsalyuk,[15] D. L. Prout,[13] V. A. Punjabi,[19] T. Pussieux,[26] G. Quéméner,[33] R. D. Ransome,[25] O. Ravel,[1] Y. Roblin,[1] R. Roche,[6] P. G. Roos,[16] D. Rowntree,[17] G.A. Rutledge,[33] P. M. Rutt,[30] A. Saha,[30] T. Saito,[31] A. J. Sarty,[6] A. Serdarevic-Offermann,[24] T. P. Smith,[18] A. Soldi,[20] P. Sorokin,[15] P. Souder,[28] R. Suleiman,[13] J. A. Templon,[7] T. Terasawa,[31] L. Todor,[21] H. Tsubota,[31] H. Ueno,[34] P. E. Ulmer,[21] G.M. Urciuoli,[12] P. Vernin,[26] S. van Verst,[17] B. Vlahovic,[20] H. Voskanyan,[35] J. W. Watson,[13] L. B. Weinstein,[21] K. Wijesooriya,[33] R. Wilson,[9] B. Wojtsekhowski,[30] D. G. Zainea,[24] V. Zeps,[14] J. Zhao,[17] Z. -L. Zhou[17]

(The Jefferson Lab Hall A Collaboration)

[1] *Université Blaise Pascal/IN2P3, F-63177 Aubière, France*
[2] *California State University, Los Angeles, California, 90032*
[3] *Institut des Sciences Nucléaires, F-38026 Grenoble, France*
[4] *Duke University, Durham, North Carolina, 27706*
[5] *Florida International University, Miami, Florida, 33199*
[6] *Florida State University, Tallahassee, Florida, 32306*
[7] *University of Georgia, Athens, Georgia, 30602*
[8] *Hampton University, Hampton, Virginia, 23668*
[9] *Harvard University, Cambridge, Massachusetts, 02138*
[10] *INFN, Sezione di Bari and University of Bari, I-70126 Bari, Italy*
[11] *INFN, Sezione di Lecce, I-73100 Lecce, Italy*
[12] *INFN, Sezione Sanitá and Istituto Superiore di Sanitá, Laboratorio di Fisica, I-00161 Rome, Italy*
[13] *Kent State University, Kent, Ohio, 44242*
[14] *University of Kentucky, Lexington, Kentucky, 40506*
[15] *Kharkov Institute of Physics and Technology, Kharkov 310108, Ukraine*
[16] *University of Maryland, College Park, Maryland, 20742*
[17] *Massachusetts Institute of Technology, Cambridge, Massachusetts, 02139*
[18] *University of New Hampshire, Durham, New Hampshire, 03824*
[19] *Norfolk State University, Norfolk, Virginia, 23504*
[20] *North Carolina Central University, Durham, North Carolina, 27707*
[21] *Old Dominion University, Norfolk, Virginia, 23529*
[22] *Institut de Physique Nucléaire, F-91406 Orsay, France*
[23] *Princeton University, Princeton, New Jersey, 08544*
[24] *University of Regina, Regina, Saskatchewan, Canada, S4S 0A2*
[25] *Rutgers, The State University of New Jersey, Piscataway, New Jersey, 08854*
[26] *CEA Saclay, F-91191 Gif-sur-Yvette, France*
[27] *State University of New York at Stony Brook, Stony Brook, New York, 11794*
[28] *Syracuse University, Syracuse, New York, 13244*
[29] *Temple University, Philadelphia, Pennsylvania, 19122*
[30] *Thomas Jefferson National Accelerator Facility, Newport News, Virginia, 23606*
[31] *Tohoku University, Sendai 980, Japan*
[32] *University of Virginia, Charlottesville, Virginia, 22901*
[33] *College of William and Mary, Williamsburg, Virginia, 23187*
[34] *Yamagata University, Yamagata 990, Japan*
[35] *Yerevan Physics Institute, Yerevan 375036, Armenia*

(September 27, 1999)

We have measured the cross section for quasielastic $1p-$shell proton knockout in the ^{16}O(e, e′p) reaction at $\omega = 0.439$ GeV and $Q^2 = 0.8$ (GeV/c)^2 for missing momentum $P_{miss} \leq 355$ MeV/c. This was the first experiment performed in Jefferson Lab Hall A. We have extracted the response functions R_{L+TT}, R_T, R_{LT}, and the left-right asymmetry A_{LT}, for the $1p_{1/2}$ and the $1p_{3/2}$ states. The data are well described by relativistic DWIA calculations. At large P_{miss}, structure observed in A_{LT} indicates the existence of dynamical relativistic effects.

PACS numbers: 25.30.Fj, 27.20.+n

Electron scattering is a powerful probe of the nuclear electromagnetic response [1,2]. Exclusive and semi-exclusive proton knockout reactions, (e, e′p), have long been used to study single-nucleon aspects of nuclear structure and to search for non-nucleonic degrees-of-freedom. At large four-momentum transfer squared [1], Q^2, quasielastic (e, e′p) is expected to be dominated by single-body interactions, hence Distorted Wave Impulse Approximation (DWIA) calculations should be more accurate than at low Q^2. Calculations [3–6] indicate that in ^{16}O(e, e′p) the longitudinal-transverse interference response function [2], R_{LT}, and the left-right asymmetry, A_{LT}, are sensitive to dynamical relativistic effects. We report structure in A_{LT} at large P_{miss} that shows for the first time clear evidence of the existence of dynamical relativistic effects in electromagnetic reactions.

^{16}O(e, e′p) $1p-$shell proton knockout experiments have been performed at Saclay [7,8], NIKHEF [9,10], and Mainz [11] at low Q^2 (less than 0.4 (GeV/c)^2) in various kinematics. These experiments measured the cross section as a function of missing momentum and have fit spectroscopic factors by comparing data to DWIA calculations. The published spectroscopic factors were between 0.5 and 0.7, but Kelly [2] showed that the data of Blomqvist *et al.* [11] suggest a significantly smaller normalization factor. Chinitz *et al.* [7] and Spaltro *et al.* [10] also extracted R_{LT}, the longitudinal-transverse interference response function at $Q^2 = 0.3$ (GeV/c)^2 and 0.2 (GeV/c)^2, respectively. Their measurements of proton knockout from the $1p_{1/2}$ state agree, but their $1p_{3/2}$ measurements disagree dramatically. DWIA calculations [6] are consistent with the data of Chinitz *et al.* [7]

This paper reports the results [12] of the first experiment [13] in Jefferson Lab Hall A [14]. In this experiment, we measured the ^{16}O(e, e′p) reaction cross section in quasielastic kinematics ($\omega = Q^2/2m_p$) at $Q^2 = 0.8$ (GeV/c)^2 and $\omega = 0.439$ GeV for $P_{miss} < 355$ MeV/c. We separated the response functions R_{L+TT}, R_T and R_{LT}, and extracted A_{LT} for $1p-$shell proton knockout.

The 100% duty factor beam current of typically 70 μA was incident on a waterfall target with three foils, each about 130 mg/cm^2 thick along the beamline [15]. We used the two Hall A High Resolution Spectrometers [14] to detect the outgoing particles. We studied the spectrometer optical properties and acceptances both before and during the experiment. The angle of the particle was determined to 0.3 mrad and its absolute momentum was measured with an accuracy $\frac{\delta p}{p} = 1.5 \times 10^{-3}$ [16–19].

The hydrogen in the H_2O target greatly simplified our normalizations and calibrations. We monitored the luminosity by continuously measuring the elastic H(e,e) cross section. We used H(e,ep) to determine the momentum transfer $|\vec{q}|$ absolutely to an accuracy of 1.5×10^{-3} and to reproduce this momentum transfer at each beam energy to a fractional accuracy of 1.5×10^{-4}.

We measured the cross section at fixed $|\vec{q}| = 992$ MeV/c at three beam energies (corresponding to three virtual photon polarizations) to separate the response functions and understand our systematic uncertainties (see Table I). The angles $\theta_{pq} = 0°, \pm 2.5°, \pm 8°, \pm 16°$, and $\pm 20°$ correspond to central missing momenta of 53, 60, 148, 280, and 345 MeV/c, respectively. Note that at $\theta_{pq} = 0°$, we had to remove events with $P_{miss} < 45$ MeV/c to eliminate contamination from H(e, ep).

For $\theta_{pq} = \pm 8°$, the values of R_{LT} and A_{LT} extracted at $E_{\rm beam} = 2.4$ GeV agree with those extracted at $E_{\rm beam} = 1.6$ GeV within one standard deviation. The overall systematic uncertainty in the response functions is about 5%. This uncertainty is dominated by the uncertainty in the H(e, e) cross section to which the data were normalized [20].

We radiatively corrected the cross section using a modified version of the code RADCOR [21]. The missing energy resolution is 0.9 MeV FWHM, which does not allow us to resolve the $(2s_{1/2}, 1d_{5/2})$ doublet located at $E_{miss} = 17.4$ MeV from the $1p_{3/2}$ state (at $E_{miss} = 18.4$ MeV). The strength of this doublet was estimated using the spectroscopic factors obtained by Leuschner *et al.* [9]

---

[1] The kinematical quantities are as follows: the scattered electron transfers momentum $\vec{q}$ and energy ω with $Q^2 = \vec{q}^2 - \omega^2$. The ejected proton has mass m_p, momentum $\vec{p}_p$, energy E_p, and kinetic energy T_p. The cross section is typically measured as a function of missing energy $E_{miss} = \omega - T_p - T_{recoil}$ and missing momentum $\vec{P}_{miss} = \vec{q} - \vec{p}_p$. The angle between the ejected proton and virtual photon is θ_{pq} and the azimuthal angle is ϕ. $\theta_{pq} > 0$ corresponds to $\phi = 180°, P_{miss} > 0$ and $\theta_p > \theta_q$. $\theta_{pq} < 0$ corresponds to $\phi = 0°$.

[2] The cross section for (e, e′p) can be written as $\frac{d^6\sigma}{d\Omega_e\, d\omega\, d\Omega_p\, dE_{miss}} = \frac{E_p p_p}{(2\pi)^3}\sigma_{\rm Mott}(V_L R_L + V_T R_T + V_{LT} R_{LT}\cos\phi + V_{TT} R_{TT}\cos 2\phi)$ where the kinematic factors $\{V_i\}$ are known and the response functions $\{R_i\}$ contain information about the nuclear charge and current densities. One can extract response functions by measuring the cross section at fixed Q^2, ω, and θ_{pq} while varying the electron scattering angle (which changes V_T) and ϕ. One can also extract the left-right asymmetry by measuring the cross section at $\phi = 0, 180°$: $A_{LT} = \frac{\sigma(\phi=0°) - \sigma(\phi=180°)}{\sigma(\phi=0°) + \sigma(\phi=180°)}$.

to be approximately 5% of the $1p_{3/2}$ strength for this kinematical region. It was not subtracted from the cross section for the $1p_{3/2}$ state.

The first relativistic calculations for (e, e′p) were performed by Picklesimer and Van Orden [3,4]. We compared our data to more recent calculations by Udias *et al.* [5,22–24] and by Kelly [6]. Both calculations use the Coulomb gauge, the NLSH bound-state wavefunction [25], the EDAIO optical potential of Cooper *et al.* [26], the cc2 prescription of the off-shell e-p cross section [27] (the use of cc1 yielded slightly poorer agreement with the data), and include the effects of electron distortion. We note that the NLSH wave function [25] yields values of binding and single-particle energies, as well as the charge radius for ^{16}O, which are in good agreement with data. Udias *et al.* solved the Dirac equation directly in configuration space, whereas Kelly solved a relativized Schrödinger equation and used the effective momentum approximation (EMA) to incorporate spinor distortion into an effective current operator based on that of Hedayati-Poor *et al.* [28]. The results for the $1p_{3/2}$ state from both calculations include an incoherent contribution from the positive-parity contaminants as parametrized by Leuschner *et al.* [9].

Figure 1 shows the cross section as a function of missing momentum at $E_{\rm beam} = 2.4$ GeV. The calculations of Udias *et al.* and of Kelly are in very good agreement with the data. This agreement is attributed to the quality of the bound-state wave function used. The spectroscopic factors are 0.73 and 0.72 for the $1p_{1/2}$ state and 0.71 and 0.67 for the $1p_{3/2}$ state for the calculations of Udias *et al.* and Kelly, respectively.

We extracted A_{LT} from the measured cross sections (see Figure 2). Note the large change in the slope of A_{LT} at $P_{miss} \approx 300$ MeV/c. This change in slope is reproduced, at least qualitatively, by both calculations, although only when they include distortions of the lower components of the Dirac spinors. This is a clear indication of the existance of dynamical relativistic effects. As can be seen in the calculations of Udias *et al.* in Figure 2, distortion of the bound-state spinors is more important than that of the ejectile spinors, although both are needed. Kelly also sees an effect due to distortion of the bound-state spinors, but due to approximations he makes, his calculations are not as accurate for $P_{miss} > 275$ MeV/c [6].

We also extracted the response functions R_{L+TT}, R_{LT}, and R_T. Since we measured the cross sections in perpendicular kinematics, we could not isolate the longitudinal response function R_L. Instead, we extracted the combination $R_{L+TT} = R_L + \frac{V_{TT}}{V_L} R_{TT}$. Both Kelly and Udias calculate the term $\frac{V_{TT}}{V_L} R_{TT}$ to be small ($< 10\% R_L$) for these kinematics. Figure 3 shows the response functions and calculations. Again, the calculations are in good agreement with the data. Moreover, neither calculation includes any two-body currents, suggesting that such currents are unimportant at this Q^2.

In summary, we have measured ^{16}O(e, e′p) $1p$−shell proton knockout in a previously inaccessible region of momentum transfer. These measurements included cross sections, the left-right asymmetry A_{LT}, and the response functions R_{L+TT}, R_{LT} and R_T for missing momentum up to approximately 350 MeV/c. The cross section, asymmetry, and response functions are reproduced very well by modern relativistic DWIA calculations without the addition of any two-body currents. Structure in A_{LT} is observed at higher missing momentum which is due to dynamic enhancement of the lower components of Dirac spinors.

We acknowledge the outstanding support of the staff of the Accelerator and Physics Divisions of Jefferson Laboratory that made this experiment successful. We thank Drs. T.W. Donnelly, J.M. Udias, and J.W. Van Orden for providing their theoretical calculations and for valuable discussions. We also thank Dr. J.M. Udias for providing us with the NLSH bound-state wave functions. This work was supported in part by the U.S. Department of Energy, the National Science Foundation, the Italian Instituto Nazionale di Fisica Nucleare (INFN), the French Atomic Energy Commission and National Center of Scientific Research, and the Natural Sciences and Engineering Research Council of Canada.

[1] S. Frullani and J. Mougey, Adv. Nucl. Phys. **14**, 1 (1984).
[2] J.J. Kelly, Adv. Nucl. Phys. **23**, 75 (1996); Phys. Rev. **C56**, 2672 (1997).
[3] A. Picklesimer, J.W. Van Orden, and S.J. Wallace, Phys. Rev. **C32**, 1312 (1985).
[4] A. Picklesimer and J.W. Van Orden, Phys. Rev. **C40**, 290 (1989).
[5] J.M. Udias *et al.*, LANL Preprint Library, nucl-th/9905030.
[6] J.J. Kelly, Phys. Rev. **C60**, 044609 (1999). The calculations were revised using the NLSH wave functions.
[7] L. Chinitz *et al.*, Phys. Rev. Lett. **67**, 568 (1991).
[8] M. Bernheim *et al.*, Nucl. Phys. **A375**, 381 (1982).
[9] M. Leuschner *et al.*, Phys. Rev. **C49**, 955 (1994).
[10] C.M. Spaltro *et al.*, Phys. Rev. **C48**, 2385 (1993).
[11] K.I. Blomqvist *et al.*, Phys. Lett. **B344**, 85 (1995).
[12] www.jlab.org/~fissum/e89003.html.
[13] A. Saha, W. Bertozzi, R.W. Lourie, and L.B. Weinstein, Jefferson Laboratory Proposal 89-003, 1989; K.G. Fissum *et al.*, MIT-LNS Internal Report #02, 1997.
[14] www.jlab.org/Hall-A/equipment/HRS.html.
[15] F. Garibaldi *et al.*, Nucl. Instrum. Meth. **A314**, 1 (1992).
[16] J. Gao, Ph.D. thesis, MIT, 1999 (unpublished).
[17] J. Gao *et al.*, MIT-LNS Internal Report #04, 1998.
[18] N. Liyanage *et al.*, MIT-LNS Internal Report #05, 1998.

[19] M. Liang, 'Survey Summary Report', Jefferson Lab Technical Notes JLAB-TN-99029.
[20] G.G. Simon *et al.*, Nucl. Phys. **A333**, 381 (1980); L.E. Price *et al.*, Phys. Rev **D4**, 45 (1971).
[21] E. Quint, Ph.D. thesis, NIKHEF 1988 (unpublished); R. Florizone, Ph.D. thesis, MIT 1999 (unpublished).
[22] J.M. Udias *et al.*, Phys. Rev. **C48**, 2731 (1993).
[23] J.M. Udias *et al.*, Phys. Rev. **C51**, 3246 (1995).
[24] J.M. Udias, private communication.
[25] M.M. Sharma, M.A. Nagarajan, and P. Ring, Phys. Lett. **B312**, 377 (1993).
[26] E.D. Cooper, S. Hama, B.C. Clark, and R.L. Mercer, Phys. Rev. **C47**, 297 (1993).
[27] T. De Forest, Nucl. Phys. **A392**, 232 (1983).
[28] M. Hedayati-Poor, J.I. Johansson, and H.S. Sherif, Phys. Rev. **C51**, 2044 (1995).

$E_{\rm beam}$ (GeV)	θ_e (°)	θ_{pq} (°)
0.843	100.7	0, 8, 16
1.643	37.2	0, ±8
2.442	23.4	0, ±2.5, ±8, ±16, ±20

TABLE I. Experimental Kinematics



FIG. 1. Measured cross sections and DWIA calculations at $E_{\rm beam} = 2.4$ GeV. The solid line is the Udias *et al.* calculation [5,24], and the dashed line is the Kelly calculation [6]. The $1p_{1/2}$ state cross section and calculations have been multiplied by a factor of 20.



FIG. 2. Measured left-right asymmetry A_{LT} and DWIA calculations at $E_{\rm beam} = 2.4$ GeV. The dashed line is the Kelly calculation [6]. The other curves are from Udias *et al.* [5,24]. The solid line is the fully relativistic calculation, the close dotted line is the calculation with only the bound-state spinor distortion included, the wide dotted line is the calculation with only the scattered-state spinor distortion included, and the dot-dashed line is the calculation without spinor distortion included.



FIG. 3. Measured R_{L+TT}, R_{LT}, R_T and DWIA calculations. The solid line is the Udias *et al.* calculation [5,24] and the dashed line is the Kelly calculation [6]. The data beyond 250 MeV/c missing momentum are expanded for clarity.