

LHRS Analysis for d_2^n

The Elastic Radiative Tail and Correcting for Background Processes

D. Flay

6/23/11

Outline

1 Cross Sections

- Radiative Corrections: The Elastic Radiative Tail
- Correcting for Background Processes

2 Summary

The Elastic Radiative Tail (1)


Discussion and Contributions

- The first step in the radiative corrections procedure is to subtract off the elastic tail from both ^3He and N
- The elastic tail (calculated in the fortran code [rosetail](#)) contains the following corrections:
 - 1 **Internal**: photon radiation before and/or after the interaction at the vertex
 - 2 **External**: straggling in the target (ionization and bremsstrahlung) and multiple-photon radiation

The Elastic Radiative Tail (2)

Testing the Code


Unpolarized Elastic ep Scattering Tail



The Elastic Radiative Tail (3)

^3He Elastic Tail


^3He 4-pass Elastic Radiative Tail



The Elastic Radiative Tail (4)

^3He Elastic Tail


^3He 5-pass Elastic Radiative Tail



The Elastic Radiative Tail (5)

Comparison to Raw and Experimental Cross Section


^3He Cross Section ($E = 4.73 \text{ GeV}, \theta = 45^\circ$)



The Elastic Radiative Tail (6)

Comparison to Raw and Experimental Cross Section

^3He Cross Section ($E = 5.89 \text{ GeV}, \theta = 45^\circ$)



Correcting for Background Processes (1)

Current Method

- Currently, we correct for e^- events coming from N_2 and those corresponding to $\gamma \rightarrow e^+e^-$ by subtracting off their cross sections:

$$\sigma_{\text{exp}} = \sigma_{\text{raw}} - \sigma_{\text{dil}} - \sigma_{e^+}$$

- σ_{raw} and σ_{dil} are the cross sections obtained on the production cell and the reference cell in negative polarity, respectively
- σ_{e^+} is obtained from the production cell in positive polarity mode.
- We also have $\sigma_{\text{dil}}^{\text{pos.}}$, obtained from the reference cell in positive polarity mode.
 - Shouldn't we also subtract this term off too?

Correcting for Background Processes (2)

Thinking in Terms of Yields

- Consider the yield for some process:

$$Y = \frac{N}{(Q/e)LT\varepsilon}$$

- The signal we detect is: $Y_{e^-} = Y_{e^-}^p + Y_{e^-}^b$
 - p = pure signal
 - b = background signal
 - $Y_{e^-}^b = Y_{e^-}^N$
 - To correct for the events scattering from N, one can subtract off $Y_{e^-}^N$
 - This is what we do for the yield from the reference cell (to remove events scattering from nitrogen)
- For the positron data, we have a similar situation:

$$Y_{e^+} = Y_{e^+}^p + Y_{e^+}^b$$

$$Y_{e^+}^b = Y_{e^+}^N$$

Correcting for Background Processes (3)

Thinking in Terms of Yields

- To determine the yield of pair-produced e^+ events that scatter from ^3He :

$$Y_{e^+}^p = Y_{e^+} - Y_{e^+}^N$$

- Therefore, the full correction should be:

$$Y_{e^-}^p = (Y_{e^-} - Y_{e^-}^N) - (Y_{e^+} - Y_{e^+}^N)$$

Summary

- Cross Section:
 - ^3He Elastic tail has been determined for 4- and 5-pass data
 - Very small contribution ($< 1\%$) for all p_0
 - Possible refined method to correct the e^- yield and effectively the cross section

What's Next?

- Acceptance:
 - Determine momentum dependence
- Cross Section:
 - Radiative Corrections:
 - N_2 elastic tail
 - Q.E., dip, DIS and $W = 1500, 1700$ MeV contributions