

0.1 E06-014

Precision Measurement of d_2^n : Probing the Lorentz Color Force

S. Choi, X. Jiang, Z.-E. Meziani, B. Sawatzky, spokespersons,
and

the d_2^n and Hall A Collaborations.

contributed by M. Posik, L. El Fassi, D. Flay, D. Parno, and Y. Zhang.

0.1.1 The Experiment

Experiment E06-014 ran in Hall A from February 7 to March 17, 2009, at two production beam energies of 4.73 and 5.89 GeV, on a polarized ^3He target. The experiment probed the resonance and deep inelastic valence quark regions, which corresponded to the ranges $0.2 \leq x \leq 0.7$ and $2 \text{ GeV}^2 \leq Q^2 \leq 6 \text{ GeV}^2$. Figure 1 shows the coverage in Q^2 and x , as well as the invariant mass of 2 GeV, which separates the deep inelastic and resonance regions. The LHRS and BigBite detector packages were each oriented at 45° relative to the beam line, with each of the detectors independent of one another, acting as its own single arm experiment. The LHRS was used to measure the unpolarized cross-section, while the BigBite measured the double-spin asymmetries in scattering between longitudinally polarized electron beam and a longitudinally and transversely polarized ^3He target.

E06-014 also served as the commissary experiment for a gas Čerenkov detector, which was installed into the the BigBite stack, as well as a new photon detector and integrating data-acquisition system for the Compton polarimeter [1, 15].




Figure 1: Q^2 vs x -Bjorken for 4.73 GeV dataset in red, 5.89 GeV dataset in blue and $W=2$ GeV is the black dashed line.

The primary goal of E06-014 is the measurement of the quantity d_2^n . The neutron d_2 is a probe of the strong force that is formed by taking the second moment of a linear combination of the polarized structure functions g_1 and g_2 :

$$d_2^n(Q^2) = \int_0^1 x^2 [2g_1^n(x, Q^2) + 3g_2^n(x, Q^2)] dx. \quad (1)$$

At low Q^2 , where the virtual photon wavelength is larger than the nucleon, d_2 can be associated with spin polarizabilities within the nucleon [3, 4]. However, at larger Q^2 , it is more appropriate to interpret d_2 as the average transverse Lorentz color force acting on a quark after being hit by a virtual photon [3, 5].

In addition to gaining insight into the nature of the color force, the precision measurement of d_2^n will also result in a bench mark test for lattice QCD predictions.

E06-014 measured d_2^n by combining the unpolarized cross-section, σ_0 from the LHRS, as well as the, parallel and perpendicular asymmetries, A_{\parallel} and A_{\perp} from BigBite. The asymmetries are defined through the counting rates of each spin orientation as:

$$A_{\parallel} = \frac{N^{\downarrow\uparrow} - N^{\uparrow\uparrow}}{N^{\downarrow\uparrow} + N^{\uparrow\uparrow}} \quad \text{and} \quad A_{\perp} = \frac{N^{\downarrow\Rightarrow} - N^{\uparrow\Rightarrow}}{N^{\downarrow\Rightarrow} + N^{\uparrow\Rightarrow}},$$

where single arrows represent the electron helicity direction, and double arrows are represent the target polarization direction. By combining these independently measured quantities, d_2^n is expressed exclusively through experimental quantities as:

$$d_2^n = \int_0^1 \frac{MQ^2}{4\alpha^2} \frac{x^2 y^2}{(1-y)(2-y)} \sigma_0 \left[\left(3 \frac{1+(1-y)\cos\theta}{(1-y)\sin\theta} + \frac{4}{y} \tan(\theta/2) \right) A_{\perp} + \left(\frac{4}{y} - 3 \right) A_{\parallel} \right] dx, \quad (2)$$

where $x = Q^2/2M\nu$ (the Bjorken x variable), $\nu = E - E'$ (the energy transfer from electron to target), θ (the scattering angle of the electron), and $y = \nu/E$ (the fractional energy transfer from electron to target). The advantage of measuring d_2^n in terms of experimental quantities, is that it allowed the allotted beam time to be divided between measuring A_{\parallel} and A_{\perp} in such a way that the error on d_2^n itself was minimized, rather than the spin structure functions g_1 and g_2 . The precision measurement of d_2^n ($\sim Q^2 = 3\text{GeV}$) is expected to result in a fourfold improvement on the world data shown in Figure 2 [6], in advance of an approved 12 GeV experiment in Hall C that will extend the precision measurement of d_2^n to higher kinematic ranges [7].




Figure 2: World data for d_2^n and several model calculations. Along with E06-014 projected d_2^n error.

In addition to the primary goal of E06-014, the data collected can also be used to measure the longitudinal virtual photon-nucleon asymmetry for the neutron, A_1^n . The virtual photon-nucleon scattering cross section can be separated into two helicity dependent cross sections, $\sigma_{1/2}$ and $\sigma_{3/2}$. The subscripts 1/2(3/2) gives the projection of the total spin along the virtual photon's momentum direction, corresponding to anti-parallel (parallel). A_1 can then be defined as:

$$A_1(x, Q^2) \equiv \frac{\sigma_{1/2} - \sigma_{3/2}}{\sigma_{1/2} + \sigma_{3/2}} \approx \frac{g_1(x, Q^2)}{F_1(x, Q^2)} \quad \text{for high } Q^2. \quad (3)$$

We may also express A_1 in terms of the parallel and perpendicular asymmetries, A_{\parallel} and A_{\perp} , that were measured in BigBite as:

$$A_1 = \frac{1}{D(1+\eta\xi)} A_{\parallel} - \frac{\eta}{d(1+\eta\xi)} A_{\perp} \quad (4)$$

where D is the virtual photon polarization factor and η , ξ , and d are quantities set by kinematics and by the virtual photon polarization vector [15].

Combining A_1^n data measured on an effective neutron target with A_1^p data measured on a polarized proton target, allows access to the polarized-unpolarized parton distribution function ratios $\Delta u/u$ and $\Delta d/d$. Recent results from Hall A [9] and from CLAS [10] showed a significant deviation of $\Delta d/d$ from the predictions of perturbative QCD, which have that ratio approaching 1 in the limit of $x \rightarrow 1$. As part of the 12 GeV program, two approved experiments (one in Hall A [11] and one in Hall C [12]) will extend the accuracy and x range of this measurement, but a measurement of A_1^n at E06-014's kinematics will provide valuable support (or refutation) of prior Jefferson Lab results, while producing additional input for theoretical models in advance of the coming experiments at 12 GeV.

0.1.2 Analysis Progress: Target and Beam

When performing a double spin asymmetry experiment, knowledge of the target and beam polarization is crucial. E06-014 used the standard Hall A polarized ^3He target with two holding field directions: longitudinal and transverse in plane with respect to the beam direction [1]. To extract the target polarization, ^3He number densities, EPR and NMR measurements were done.

The number density of ^3He was measured in both the pumping chamber and the target chamber. This measurement was achieved by using the fact that collisions with ^3He atoms broaden the D1 and D2 absorption lines of rubidium [17]. By measuring the width of the D1 and D2 absorption lines and subtracting 1% N_2 contribution, a measurement of ^3He number density at room temperature, n_0 , can be obtained. Since the number density changes with temperature, equations 5 and 6, were used to compute the number densities in both the pumping and target chambers, where V_{tot} is the total volume of the target cell, T is the temperature and the subscripts $pc(tc)$ refer to the pumping(target) chamber. The temperature of the chambers was measured using seven resistive thermal devices (RTDs), which were placed outside of the target and were stable within 2°C during production [15].

$$n_{pc} = n_0 \left[1 + \frac{V_{pc}}{V_{tot}} \left(\frac{T_{tc}}{T_{pc}} - 1 \right) \right]^{-1} \quad (5)$$

$$n_{tc} = n_0 \left[1 + \frac{V_{tc}}{V_{tot}} \left(\frac{T_{pc}}{T_{tc}} - 1 \right) \right]^{-1} \quad (6)$$

The ^3He number densities for the E06-014 target cell Samantha as a function of run number can be seen in Figure 3, with the average values listed in Table 1.




Figure 3: ^3He densities as a function of BigBite run number [15, 16]

During E06-014 EPR measurements were taken every several days, while NMR measurements were taken every four hours. During EPR measurements, the frequency shift of potassium level transitions in

Chamber	^3He Density (amg)
Pumping	6.93 ± 0.19
Target	10.81 ± 0.29

Table 1: Average ^3He densities in target cell [13, 15]

the presence of polarized ^3He were measured, this frequency shift $\Delta\nu_{EPR}$, can be related to the target polarization, $P_{^3\text{He}}$:

$$\Delta\nu_{EPR} = \frac{4\mu_0}{3} \frac{d\nu_{EPR}}{dB} \kappa_0 \mu_{^3\text{He}} n_{pc} P_{^3\text{He}}. \quad (7)$$

where μ_0 is the vacuum permeability, $\mu_{^3\text{He}}$ is the magnetic moment, $\frac{d\nu_{EPR}}{dB}$ is the derivative of the EPR frequency with respect to the magnetic field, κ_0 is the enhancement factor and n_{pc} is the pumping chamber number density. During EPR measurements, a NMR measurement was also done. This allows a comparison between the EPR polarization, $P_{^3\text{He}}$, and the measured NMR amplitude, h : This conversion factor allows NMR measurements to be converted into an absolute ^3He polarization, and is defined as:

$$c' = \frac{P_{^3\text{He}}}{h}. \quad (8)$$

In addition to obtaining the conversion factor c' from using EPR and NMR measurements, it can also be calculated by performing a calibration on a water target. The final NMR polarization, is then computed by taking the weighted average of the polarization computed from the EPR and water calibrations. Although the water calibration still needs to be done in order to obtain the final target polarization, the EPR measurements can be used to obtain a preliminary target polarization. The average conversion factor c' for all EPR measurements in each target polarization direction was then computed and applied to the NMR measurements Table 2. A linear interpolation was then done in order to obtain the polarization on a run by run basis. This procedure resulted in a combined systematic and statistical error of 4.9% Figure 4 [13, 15].

Polarization Direction	c' (%/mV)
Longitudinal	2.84 ± 0.14
Transverse	1.77 ± 0.09

Table 2: EPR-NMR conversion factors c' [14, 15]

In addition to the target polarization, the beam polarization also needs to be know. E06-014 used a polarized electron beam at energies of 4.73 and 5.89 GeV. The polarization of the electron beam was measured independently through Compton and Møller scattering. During the running of E06-014, there were several Møller measurements performed, while Compton data was taken continuously through out the experiment. Figure 5 shows the beam polarization as a function of BigBite run number for the Møller and Compton results. The beam polarization data was split into four run sets and the average polarization for each run period was then computed by taking into account both the Compton and Møller data. The final beam polarizations can be seen in Table 3 [15].

Run Set	Beam Energy (GeV)	P_e from Compton	P_e from Møller	Combined P_e
1	5.90	0.726 ± 0.018	0.745 ± 0.015	0.737 ± 0.012
2	4.74	0.210 ± 0.011	-	0.210 ± 0.011
3	5.90	0.787 ± 0.020	0.797 ± 0.016	0.793 ± 0.012
4	4.74	0.623 ± 0.016	0.628 ± 0.012	0.626 ± 0.010

Table 3: Final beam polarization for E06-014. For run set 2 there was no Møller measurement. [15]




Figure 4: Preliminary target polarization based on linear interpolation of NMR polarization measurements calibrated using EPR results. Red lines show spin transitions between transverse and longitudinal orientations. Blue lines show 180° rotations of spin orientation within a polarization plane. The labels at the top of the graph give polarization direction during that time period: $L_+ = 0^\circ$, $T_- = 90^\circ$ and $T_+ = 270^\circ$ [15, 16].




Figure 5: Final electron beam polarization from Møller and Compton measurements for E06-014. Note there was no Møller measurements for the second run set [15].

0.1.3 Analysis Progress: LHRS

The LHRS was used to measure the total unpolarized cross-section, which will multiply the measured asymmetries in BigBite which is needed to compute d_2^u . In order to measure the unpolarized cross-section, the LHRS hardware must be calibrated and PID cuts applied in order to select electrons from the data [1]. The total unpolarized cross-section is given as:

$$\sigma_0 = \frac{N \cdot ps \cdot e}{\epsilon \cdot w \cdot Q \cdot LT \cdot \rho \cdot \Delta Z \Delta \theta \Delta \phi \Delta E'} \quad (9)$$

where N is the number of events counted, ps is the pre-scale factor, ϵ is the detector efficiencies, Q is the total charge on target, LT is the trigger live time, ρ is the target density, ΔZ is the effective target length as seen by the LHRS, $\Delta \theta$ is the vertical (dispersive) direction, ϕ is the horizontal (transverse) direction, $\Delta E'$ is the energy width seen by the LHRS, and w is an acceptance weight factor that is determined from Monte Carlo simulation that signifies the percentage of events (generated at the target) that reach the LHRS focal plane. In order to determine the weight factor w , a Single Arm Monte Carlo (SAMC) simulation was utilized. SAMC determines how the geometrical acceptance of the LHRS deviates from an ideal square distribution. This is done by having SAMC generate a normal distribution in θ, ϕ and $\delta p/p$. Then with the use of John

LeRose transport matrices [18], the events are propagated to the focal plane of the LHRs. As the particle encounters each magnet aperture in the LHRs, a check is performed to see if the particle passes through the aperture until it reaches the focal plane. For all events that make it to the focal plane, the event is then reconstructed at the target. The ratio of events that make it to the focal plane to the generated events forms the acceptance weight factor w . In Figure 6, we see the results from SAMC to actual production data for the target length seen by the LHRs, the vertical and horizontal angular distributions and $\delta p/p$. The acceptance weight is typically on the order of 0.99.




Figure 6: Comparison of events that passed in SAMC to E06-014 production data from LHRs.

The raw ${}^3\text{He}$ cross-section that was measured in the LHRs, σ_{raw} , contains contributions from positrons and nitrogen that must be corrected for in order to have a pure electron cross-section. In order to remove these contributions from σ_{raw} , several runs were taken with the LHRs in positive polarity mode, in which positrons were detected. From these runs a positron cross-section, σ_{e+} was measured. To obtain the amount of nitrogen that is present in the ${}^3\text{He}$ runs, electrons were scattered off of a nitrogen target, and a nitrogen cross-section, σ_{N_2} was measured. In order to use σ_{N_2} to correct σ_{raw} , the differences in the ${}^3\text{He}$ and N_2 targets must be accounted for. This is corrected for by taking the ratio of the target densities:

$$\sigma_{N_2}^{dil} = \frac{n_{N_2}}{n_{N_2} + n_{3He}} \sigma_{N_2}, \quad (10)$$

where n_{N_2} is the number density of the N_2 target cell and n_{3He} is the number density of the ${}^3\text{He}$ target. σ_{raw} can now be corrected for positron and nitrogen contributions by:

$$\sigma_{exp} = \sigma_{raw} - \sigma_{e+} - \sigma_{N_2}^{dil} \quad (11)$$

Figure 7, 8 show the the 4.74 and 5.89 GeV raw, positron, nitrogen and corrected cross-sections measured in the LHRs as a function of x . With the preliminary 4.74 and 5.89 GeV cross-sections completed, analysis is now being done on radiative corrections and fitting the cross-section data.

0.1.4 Analysis Progress: BigBite

E06-014 used the BigBite detector package to measure the double spin asymmetry between longitudinally polarized electrons and a longitudinally and transversely polarized ${}^3\text{He}$ target. The BigBite detector calibrations and data quality checks have been completed [1, 15]. Preliminary asymmetry analysis at a beam energy of 4.74 GeV has also been completed. The longitudinal and transverse asymmetries were computed using Equation 2. The kinematic range has been divided ($0.15 \leq x \leq 1.0$) into 17 equally space x bins.




Figure 7: LHRs 4.74 GeV ^3He cross-sections as a function of x . Error bars on the cross-sections are statistical only, and have not been corrected for radiative corrections.




Figure 8: LHRs 5.89 GeV ^3He cross-sections as a function of x . Error bars on the cross-sections are statistical only, and have not been corrected for radiative corrections.

As can be seen from Figure 1, this data set is in the deep-inelastic scattering (DIS) regime ($W < 2$) for $x \leq 0.55$; above this value, the data are in the resonance region.

Since our ^3He target also has a small percentage of N_2 present, the unpolarized N_2 gas will tend to dilute the measured asymmetries. In order to correct for this, counting rates from a pure N_2 target was also measured. Comparing the N_2 target counting rates to the ^3He production cell scattering rates, a dilution factor can be formed and applied to the measured asymmetry. The dilution factor is given by Equation (12).

$$D_{N_2} = 1 - \frac{\Sigma_{N_2}(N_2)}{\Sigma_{total}(^3\text{He})} \frac{Q(^3\text{He})\rho_{N_2}(^3\text{He})}{Q(N_2)\rho_{N_2}(N_2)} \quad (12)$$

Where Σ_{N_2} is that total scattering counts detected during the N_2 target run, Σ_{total} is the total scattering counts detected during a ^3He target run. $Q(N_2)$, $Q(^3\text{He})$ is the total charge deposited on the two targets and $\rho_{N_2}(N_2)$, $\rho_{N_2}(^3\text{He})$ are the nitrogen number densities present in the two targets. The dilution factor was computed at each of the 17 x bins Figure 9.

In addition to correcting for nitrogen dilution, corrections for target and beam polarizations must also be applied since the target and beam were not fully polarized. This is accounted for by dividing the asymmetries by the measured target and beam polarizations, P_t and P_b , found in Figure 4 for the last run set, and Table 3. The physics asymmetries are then given as:

$$A_{\parallel}^{phys} = \frac{1}{P_b P_t D_{N_2}} A_{\parallel}, A_{\perp}^{phys} = \frac{1}{P_b P_t D_{N_2} \cos(\phi)} A_{\perp} \quad (13)$$

,where ϕ is the vertical angle. Figure 10 shows the physics asymmetries on ^3He at an electron beam energy of 4.74 GeV as a function of x . These asymmetries can also be used to from longitudinal virtual photon-nucleon asymmetry on ^3He , $A_1^{^3\text{He}}$ via Equation 4. Preliminary E06-014 A_1 asymmetry measurements on ^3He are in good agreement with previous A_1 asymmetry measurements, as can be seen in Figure 11. E06-014's largest sources of systematic error on the 4.74 GeV data set are due to the target polarization measurement, from the contamination of the DIS sample by pair-production processes, and from errors in momentum assignment. Completing and finalizing the target water calibration will substantially reduce the systematic error. A complete study of backgrounds must await analysis of the larger 5.89 GeV data set, during which time trigger thresholds were lower and correspondingly background rates were higher. Calibrations and data quality checks have just been completed on the 5.89 GeV data set, with asymmetry and background studies to follow shortly. The extraction of the neutron asymmetry A_1^n is also underway. However, the extraction of the neutron asymmetry A_1^n is model-dependent. Previous experiments [9] have used Bissey *et al.*'s complete




Figure 9: The N_2 dilution factor is plotted vs x at a beam energy of 4.74 GeV. Outer error bars show combined systematic and statistical errors. The inner error bars show only statistical errors. In some bins, the statistical error is too small to be seen on the graph.

model in the DIS regime [20]. However, E06-014's 4.74 GeV data set spans both the DIS and resonance regions. A consistent treatment of both DIS and resonance data requires careful consideration of structure-function smearing [21]. We are working with W. Melnitchouk to extract neutron asymmetries across our entire kinematic range.




Figure 10: A_{\parallel} and A_{\perp} on ${}^3\text{He}$ are plotted vs x at a beam energy of 4.74 GeV [15].




Figure 11: Preliminary E06-014 measurement of $A_1^{3\text{He}}$ at 4.74 GeV, compared to ^3He data from E142 at SLAC [19] and E99-117 in Hall A [9] [15].

References

- [1] D. Parno, <http://hallaweb.jlab.org/experiment/E06-014/talks/index.html>, E06014 Analysis Status Report (12/2010)
- [2] S. Wandzura and F. Wilczek, Phys. Lett. **B 72**, 195 (1977).
- [3] M. Burkardt, hep-ph/0905.4079v1 (2009).
- [4] E. Stein *et al.*, Phys. Lett. B **343**, 369 (1995).
- [5] M. Burkardt, AIP Conference Proceedings **1149**, 62 (2008), hep-ph/0902.0163v1.
- [6] S. Choi, Z.-E. Meziani, X. Jiang, B. Sawatzky, *et al.*, Jefferson Lab PAC PR-06-014 (2005).
- [7] T. Averett, W. Korsch, Z.-E. Meziani, B. Sawatzky *et al.*, Jefferson Lab PAC E1206121 (2010).
- [8] X. Zheng *et al.*, Phys. Rev. Lett. **92**, 012004 (2004).
- [9] X. Zheng *et al.*, Phys. Rev. C **70**, 065207 (2004).
- [10] K. V. Dharmawardane *et al.*, Phys. Lett. B **641**, 11 (2006).
- [11] G. Cates, N. Liyanage, Z.-E. Meziani, G. Rosner, B. Wojtsekhowski, X. Zheng, *et al.*, Jefferson Lab PAC E1206122 (2006).
- [12] G. Cates, J.-P. Chen, Z.-E. Meziani, X. Zheng *et al.*, Jefferson Lab PAC E1210101 (2010).
- [13] Y. Zhang, Target analysis for E06-014. http://jlab.org/yawei/d2n/targ_ana.html
- [14] Y. Zhang, Electron paramagnetic resonance (EPR) project summary. Technical report, Hall A Collaboration (2011), <http://www.jlab.org/yawei/weeklymeeting/110209/epr.pdf>
- [15] D. Parno, PhD thesis, Carnegie Mellon University (2011).
- [16] Y. Zhang, Private communication
- [17] I. Kominis, Measurement of the neutron (^3He) spin structure at low Q^2 and the extended Gerasimov-Drell-Hearn sum rule. Ph.D. thesis, Princeton University (2001)

- [18] J. J. Leroose, Transfer Functions for Hall A Spectrometers, <http://hallaweb.jlab.org/news/minutes/tranferfuncs.html>
- [19] P. L. Anthony *et al.*, Phys. Rev. D **54**, 6620 (1996).
- [20] F. Bissey *et al.*, Phys. Rev. C **65**, 064317 (2002).
- [21] , S. A. Kulagin and W. Melnitchouk, Phys. Rev. C **78**, 065203 (2008).